



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 25, 2024 – 03:38 PM EST

PDB ID : 6U26
Title : PCSK9 in complex with compound 16
Authors : Orth, P.
Deposited on : 2019-08-19
Resolution : 1.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	 13% 87%
1	B	707	 9% 63% 32%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4633 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	92	739	474	133	130	2	0	0	0
1	B	479	3498	2161	641	666	30	0	0	0

There are 94 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	474	ILE	VAL	engineered mutation	UNP Q8NBP7
A	670	GLU	GLY	engineered mutation	UNP Q8NBP7
A	693	LYS	-	expression tag	UNP Q8NBP7
A	694	GLY	-	expression tag	UNP Q8NBP7
A	695	ASN	-	expression tag	UNP Q8NBP7
A	696	SER	-	expression tag	UNP Q8NBP7
A	697	ALA	-	expression tag	UNP Q8NBP7
A	698	ASP	-	expression tag	UNP Q8NBP7
A	699	ILE	-	expression tag	UNP Q8NBP7
A	700	GLN	-	expression tag	UNP Q8NBP7
A	701	HIS	-	expression tag	UNP Q8NBP7
A	702	SER	-	expression tag	UNP Q8NBP7
A	703	GLY	-	expression tag	UNP Q8NBP7
A	704	GLY	-	expression tag	UNP Q8NBP7
A	705	ARG	-	expression tag	UNP Q8NBP7
A	706	SER	-	expression tag	UNP Q8NBP7
A	707	SER	-	expression tag	UNP Q8NBP7
A	708	LEU	-	expression tag	UNP Q8NBP7
A	709	GLU	-	expression tag	UNP Q8NBP7
A	710	GLY	-	expression tag	UNP Q8NBP7
A	711	PRO	-	expression tag	UNP Q8NBP7
A	712	ARG	-	expression tag	UNP Q8NBP7
A	713	PHE	-	expression tag	UNP Q8NBP7
A	714	GLU	-	expression tag	UNP Q8NBP7
A	715	GLY	-	expression tag	UNP Q8NBP7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	716	LYS	-	expression tag	UNP Q8NBP7
A	717	PRO	-	expression tag	UNP Q8NBP7
A	718	ILE	-	expression tag	UNP Q8NBP7
A	719	PRO	-	expression tag	UNP Q8NBP7
A	720	ASN	-	expression tag	UNP Q8NBP7
A	721	PRO	-	expression tag	UNP Q8NBP7
A	722	LEU	-	expression tag	UNP Q8NBP7
A	723	LEU	-	expression tag	UNP Q8NBP7
A	724	GLY	-	expression tag	UNP Q8NBP7
A	725	LEU	-	expression tag	UNP Q8NBP7
A	726	ASP	-	expression tag	UNP Q8NBP7
A	727	SER	-	expression tag	UNP Q8NBP7
A	728	THR	-	expression tag	UNP Q8NBP7
A	729	ARG	-	expression tag	UNP Q8NBP7
A	730	THR	-	expression tag	UNP Q8NBP7
A	731	GLY	-	expression tag	UNP Q8NBP7
A	732	HIS	-	expression tag	UNP Q8NBP7
A	733	HIS	-	expression tag	UNP Q8NBP7
A	734	HIS	-	expression tag	UNP Q8NBP7
A	735	HIS	-	expression tag	UNP Q8NBP7
A	736	HIS	-	expression tag	UNP Q8NBP7
A	737	HIS	-	expression tag	UNP Q8NBP7
B	474	ILE	VAL	engineered mutation	UNP Q8NBP7
B	670	GLU	GLY	engineered mutation	UNP Q8NBP7
B	693	LYS	-	expression tag	UNP Q8NBP7
B	694	GLY	-	expression tag	UNP Q8NBP7
B	695	ASN	-	expression tag	UNP Q8NBP7
B	696	SER	-	expression tag	UNP Q8NBP7
B	697	ALA	-	expression tag	UNP Q8NBP7
B	698	ASP	-	expression tag	UNP Q8NBP7
B	699	ILE	-	expression tag	UNP Q8NBP7
B	700	GLN	-	expression tag	UNP Q8NBP7
B	701	HIS	-	expression tag	UNP Q8NBP7
B	702	SER	-	expression tag	UNP Q8NBP7
B	703	GLY	-	expression tag	UNP Q8NBP7
B	704	GLY	-	expression tag	UNP Q8NBP7
B	705	ARG	-	expression tag	UNP Q8NBP7
B	706	SER	-	expression tag	UNP Q8NBP7
B	707	SER	-	expression tag	UNP Q8NBP7
B	708	LEU	-	expression tag	UNP Q8NBP7
B	709	GLU	-	expression tag	UNP Q8NBP7
B	710	GLY	-	expression tag	UNP Q8NBP7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	711	PRO	-	expression tag	UNP Q8NBP7
B	712	ARG	-	expression tag	UNP Q8NBP7
B	713	PHE	-	expression tag	UNP Q8NBP7
B	714	GLU	-	expression tag	UNP Q8NBP7
B	715	GLY	-	expression tag	UNP Q8NBP7
B	716	LYS	-	expression tag	UNP Q8NBP7
B	717	PRO	-	expression tag	UNP Q8NBP7
B	718	ILE	-	expression tag	UNP Q8NBP7
B	719	PRO	-	expression tag	UNP Q8NBP7
B	720	ASN	-	expression tag	UNP Q8NBP7
B	721	PRO	-	expression tag	UNP Q8NBP7
B	722	LEU	-	expression tag	UNP Q8NBP7
B	723	LEU	-	expression tag	UNP Q8NBP7
B	724	GLY	-	expression tag	UNP Q8NBP7
B	725	LEU	-	expression tag	UNP Q8NBP7
B	726	ASP	-	expression tag	UNP Q8NBP7
B	727	SER	-	expression tag	UNP Q8NBP7
B	728	THR	-	expression tag	UNP Q8NBP7
B	729	ARG	-	expression tag	UNP Q8NBP7
B	730	THR	-	expression tag	UNP Q8NBP7
B	731	GLY	-	expression tag	UNP Q8NBP7
B	732	HIS	-	expression tag	UNP Q8NBP7
B	733	HIS	-	expression tag	UNP Q8NBP7
B	734	HIS	-	expression tag	UNP Q8NBP7
B	735	HIS	-	expression tag	UNP Q8NBP7
B	736	HIS	-	expression tag	UNP Q8NBP7
B	737	HIS	-	expression tag	UNP Q8NBP7

- Molecule 2 is 4'-{[(1R)-6-{2-[2-({N 5 -[N,N'-bis(tert-butoxycarbonyl)carbamimidoyl]-N 2 -(tert-butoxycarbonyl)-L-ornithyl}amino)ethoxy]ethoxy}-1-methyl-1-{2-oxo-2-[(1,3-thiazol-2-yl)amino]ethyl}-1,2,3,4-tetrahydroisoquinolin-7-yl]oxy}-2'-fluoro[1,1'-biphenyl]-4-carboxylic acid (three-letter code: 063) (formula: C₅₃H₆₉FN₈O₁₃S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
2	B	1	76	53	1	8	13	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	77	Total	O	0	0
			77	77		
3	B	243	Total	O	0	0
			243	243		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.68Å 70.92Å 150.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.12 – 1.53 22.12 – 1.53	Depositor EDS
% Data completeness (in resolution range)	74.5 (22.12-1.53) 74.5 (22.12-1.53)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 1.53Å)	Xtrriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.187 , 0.213 0.191 , 0.219	Depositor DCC
R_{free} test set	3848 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtrriage
Anisotropy	0.038	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4633	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 063

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/756	0.58	0/1023
1	B	0.50	0/3561	0.63	0/4844
All	All	0.50	0/4317	0.62	0/5867

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	739	0	750	1	0
1	B	3498	0	3371	15	0
2	B	76	0	0	1	0
3	A	77	0	0	1	0
3	B	243	0	0	1	0
All	All	4633	0	4121	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:185:LEU:HD11	1:B:271:ILE:HD11	1.77	0.66
1:A:138:PRO:HD2	3:A:808:HOH:O	1.97	0.62
1:B:416:ILE:HD11	3:B:1001:HOH:O	2.00	0.62
1:B:186:ASP:OD1	1:B:288:PRO:HG2	2.02	0.58
1:B:201:MET:HE2	1:B:248:ARG:HH21	1.73	0.54
1:B:584:GLN:HE21	1:B:585:PRO:HD2	1.75	0.52
1:B:490:SER:HB2	1:B:520:VAL:HG12	1.91	0.52
1:B:483:LEU:HD22	1:B:507:LEU:HD22	1.92	0.51
1:B:446:PRO:HB2	1:B:448:THR:HG23	1.94	0.50
1:B:549:ARG:HG2	1:B:589:VAL:HG22	1.97	0.47
1:B:196:ILE:HB	1:B:200:VAL:HG23	2.00	0.43
1:B:468:THR:HB	1:B:471:ALA:HB2	2.01	0.42
1:B:313:THR:HG21	1:B:328:ALA:HA	2.01	0.41
1:B:163:PRO:HD3	1:B:444:LEU:HB2	2.02	0.41
2:B:801:063:O3	2:B:801:063:S5	2.79	0.41
1:B:206:GLU:HG3	1:B:251:ARG:HD3	2.03	0.40
1:B:201:MET:HE2	1:B:248:ARG:NH2	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	90/707 (13%)	87 (97%)	3 (3%)	0	100	100
1	B	463/707 (66%)	452 (98%)	9 (2%)	2 (0%)	34	13
All	All	553/1414 (39%)	539 (98%)	12 (2%)	2 (0%)	34	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	643	HIS
1	B	280	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/571 (14%)	78 (99%)	1 (1%)	69	42
1	B	370/571 (65%)	365 (99%)	5 (1%)	67	39
All	All	449/1142 (39%)	443 (99%)	6 (1%)	69	42

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	LEU
1	B	375	CYS
1	B	448	THR
1	B	468	THR
1	B	570	ASP
1	B	623	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	464	HIS
1	B	584	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	063	B	801	-	77,80,80	1.03	6 (7%)	96,114,114	1.20	9 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	063	B	801	-	-	18/67/82/82	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	063	C55-N15	-3.60	1.30	1.37
2	B	801	063	C55-N14	3.00	1.38	1.33
2	B	801	063	C50-N49	2.40	1.38	1.33
2	B	801	063	C17-N1	2.34	1.50	1.47
2	B	801	063	C55-N13	2.20	1.38	1.34
2	B	801	063	O63-C64	-2.02	1.44	1.48

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	063	N15-C55-N13	3.51	127.77	118.08
2	B	801	063	C64-O63-C68	-3.08	118.47	121.32
2	B	801	063	C74-C71-C73	-3.04	103.19	111.16
2	B	801	063	C55-N15-C61	-2.47	119.67	125.02
2	B	801	063	O63-C68-N14	2.19	111.19	108.59
2	B	801	063	C55-N14-C68	2.15	123.58	119.66
2	B	801	063	F8-C36-C35	2.14	122.11	118.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	063	C29-N2-C28	-2.12	123.81	129.54
2	B	801	063	O6-C23-C22	-2.08	115.67	121.80

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	801	063	N14-C55-N13-C54
2	B	801	063	N15-C55-N13-C54
2	B	801	063	N13-C55-N14-C68
2	B	801	063	N15-C55-N14-C68
2	B	801	063	O56-C61-N15-C55
2	B	801	063	O62-C61-N15-C55
2	B	801	063	C50-C51-C52-C53
2	B	801	063	N14-C68-O63-C64
2	B	801	063	O69-C68-O63-C64
2	B	801	063	C65-C64-O63-C68
2	B	801	063	N16-C51-C52-C53
2	B	801	063	C67-C64-O63-C68
2	B	801	063	C66-C64-O63-C68
2	B	801	063	O9-C41-C48-N49
2	B	801	063	C51-C52-C53-C54
2	B	801	063	C39-C40-O9-C41
2	B	801	063	N1-C18-C26-C28
2	B	801	063	C27-C18-C26-C28


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	063	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	92/707 (13%)	0.07	5 (5%) 25 29	15, 21, 43, 56	0
1	B	479/707 (67%)	0.64	65 (13%) 3 2	13, 23, 59, 94	0
All	All	571/1414 (40%)	0.55	70 (12%) 4 3	13, 23, 57, 94	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	515	PHE	9.4
1	B	505	GLY	7.8
1	B	643	HIS	7.7
1	B	466	GLY	7.5
1	B	467	PRO	7.4
1	B	451	ALA	7.4
1	B	556	GLY	7.3
1	B	585	PRO	7.2
1	B	468	THR	7.1
1	B	568	VAL	6.8
1	B	220	ALA	6.6
1	B	615	ILE	6.6
1	B	642	SER	6.4
1	B	516	GLY	6.4
1	B	504	GLY	6.2
1	B	570	ASP	5.8
1	B	555	GLN	5.8
1	B	448	THR	5.7
1	B	548	THR	5.6
1	B	671	ALA	5.5
1	B	549	ARG	5.3
1	B	589	VAL	5.2
1	B	452	GLY	5.0
1	B	450	GLY	5.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	212	ASP	4.8
1	B	569	GLU	4.8
1	B	449	HIS	4.6
1	A	61	THR	4.4
1	B	470	MET	4.1
1	B	503	GLN	4.1
1	B	616	PRO	4.1
1	B	517	GLY	4.0
1	B	602	HIS	4.0
1	B	210	GLU	4.0
1	B	453	TRP	4.0
1	B	591	HIS	3.9
1	B	221	SER	3.9
1	B	514	ALA	3.9
1	B	533	ASN	3.6
1	B	530	PRO	3.4
1	B	531	GLN	3.4
1	B	553	HIS	3.3
1	B	584	GLN	3.2
1	B	659	ARG	3.1
1	B	178	SER	3.1
1	B	447	SER	3.1
1	B	587	GLN	3.1
1	B	639	PRO	3.0
1	B	604	PRO	3.0
1	B	153	SER	3.0
1	B	492	SER	3.0
1	B	644	VAL	2.9
1	B	592	ARG	2.9
1	B	534	CYS	2.8
1	A	108	LEU	2.7
1	A	85	GLU	2.7
1	B	167	ARG	2.7
1	B	204	ASP	2.7
1	B	469	ARG	2.6
1	B	677	ILE	2.6
1	B	542	ALA	2.4
1	B	658	SER	2.4
1	B	567	GLU	2.4
1	B	532	ALA	2.3
1	B	557	HIS	2.3
1	B	502	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	143	ILE	2.2
1	B	550	VAL	2.2
1	A	104	ARG	2.1
1	B	554	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	063	B	801	76/76	0.93	0.14	13,26,91,93	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.