



Full wwPDB X-ray Structure Validation Report i

Oct 4, 2023 – 09:32 PM EDT

PDB ID : 6UXQ
Title : Crystal structure of BAK core domain BH3-groove-dimer in complex with POPC and C8E4
Authors : Cowan, A.D.; Colman, P.M.; Czabotar, P.E.
Deposited on : 2019-11-07
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see references ①) were used in the production of this report:

MolProbity	: FAILED
Mogul	: 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	: 1.13
EDS	: FAILED
buster-report	: 1.1.7 (2018)
Percentile statistics	: 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	: Engh & Huber (2001)
Ideal geometry (DNA, RNA)	: Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	: 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

There are no overall percentile quality scores available for this entry.

MolProbit and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [\(i\)](#)

There are 6 unique types of molecules in this entry. The entry contains 3034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bcl-2 homologous antagonist/killer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	84	Total	C 702	N 449	O 121	S 130	2	0	5	0
1	B	80	Total	C 672	N 427	O 117	S 125	3	0	4	0
1	C	80	Total	C 664	N 426	O 116	S 120	2	0	3	0
1	D	81	Total	C 661	N 419	O 115	S 125	2	0	2	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	64	GLY	-	expression tag	UNP Q16611
A	65	PRO	-	expression tag	UNP Q16611
A	66	LEU	-	expression tag	UNP Q16611
A	67	GLY	-	expression tag	UNP Q16611
B	64	GLY	-	expression tag	UNP Q16611
B	65	PRO	-	expression tag	UNP Q16611
B	66	LEU	-	expression tag	UNP Q16611
B	67	GLY	-	expression tag	UNP Q16611
C	64	GLY	-	expression tag	UNP Q16611
C	65	PRO	-	expression tag	UNP Q16611
C	66	LEU	-	expression tag	UNP Q16611
C	67	GLY	-	expression tag	UNP Q16611
D	64	GLY	-	expression tag	UNP Q16611
D	65	PRO	-	expression tag	UNP Q16611
D	66	LEU	-	expression tag	UNP Q16611
D	67	GLY	-	expression tag	UNP Q16611

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

- Molecule 3 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (three-letter code: C8E) (formula: C₁₆H₃₄O₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 7 7	0	0
3	A	1	Total C O 11 10 1	0	0
3	A	1	Total C O 21 16 5	0	0
3	B	1	Total C O 21 16 5	0	0
3	B	1	Total C O 9 8 1	0	0
3	B	1	Total C O 21 16 5	0	0
3	C	1	Total C O 15 12 3	0	0
3	D	1	Total C O 17 12 5	0	0


- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (three-letter code: LBN)

(formula: C₄₂H₈₂NO₈P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	B	1	35	25	1	8	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	45	Total	O 45	0	0
6	B	43	Total	O 43	0	0
6	C	26	Total	O 26	0	0
6	D	32	Total	O 32	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	84.32 Å 150.09 Å 51.75 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.02 – 1.70	Depositor
% Data completeness (in resolution range)	99.2 (43.02-1.70)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	0.99 (at 1.70 Å)	Xtriage
Refinement program	PHENIX 1.15-3459	Depositor
R , R_{free}	0.179 , 0.214	Depositor
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.603	Xtriage
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.015 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.023 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
Total number of atoms	3034	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [\(i\)](#)

4.1 Standard geometry [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [\(i\)](#)

4.3.1 Protein backbone [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [\(i\)](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	C8E	A	505	-	20,20,20	0.39	0	19,19,19	0.47	0
5	LBN	B	202	-	34,34,51	1.23	3 (8%)	40,42,59	0.92	2 (5%)
3	C8E	C	200	-	14,14,20	0.40	0	13,13,19	0.34	0
2	SO4	D	202	-	4,4,4	0.19	0	6,6,6	0.27	0
2	SO4	A	502	-	4,4,4	0.13	0	6,6,6	0.11	0
2	SO4	A	501	-	4,4,4	0.19	0	6,6,6	0.26	0
4	EDO	A	506	-	3,3,3	0.48	0	2,2,2	0.32	0
4	EDO	B	207	-	3,3,3	0.41	0	2,2,2	0.51	0
3	C8E	B	203	-	8,8,20	0.26	0	7,7,19	0.52	0
3	C8E	A	503	-	6,6,20	0.27	0	5,5,19	0.43	0
3	C8E	D	201	-	16,16,20	0.41	0	15,15,19	0.35	0
2	SO4	B	204	-	4,4,4	0.14	0	6,6,6	0.09	0
3	C8E	B	205	-	20,20,20	0.39	0	19,19,19	0.44	0
3	C8E	A	504	-	10,10,20	0.37	0	9,9,19	0.45	0
4	EDO	B	206	-	3,3,3	0.35	0	2,2,2	1.10	0
3	C8E	B	201	-	20,20,20	0.40	0	19,19,19	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	C8E	A	505	-	-	12/18/18/18	-
5	LBN	B	202	-	-	8/38/38/55	-
3	C8E	C	200	-	-	6/12/12/18	-
4	EDO	A	506	-	-	0/1/1/1	-
4	EDO	B	207	-	-	0/1/1/1	-
3	C8E	B	203	-	-	4/6/6/18	-
3	C8E	A	503	-	-	4/4/4/18	-
3	C8E	D	201	-	-	8/14/14/18	-
3	C8E	B	205	-	-	10/18/18/18	-
3	C8E	A	504	-	-	4/8/8/18	-
4	EDO	B	206	-	-	1/1/1/1	-
3	C8E	B	201	-	-	9/18/18/18	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	202	LBN	O5-C25	2.85	1.41	1.33
5	B	202	LBN	O7-C34	2.79	1.42	1.34
5	B	202	LBN	O7-C2	-2.38	1.40	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	202	LBN	O7-C34-C35	3.82	119.73	111.50
5	B	202	LBN	O5-C25-C26	2.44	119.57	111.91

There are no chirality outliers.

All (66) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	202	LBN	O7-C2-C3-O5
3	A	505	C8E	C17-C16-O15-C14
3	D	201	C8E	O15-C16-C17-O18
3	B	201	C8E	O15-C16-C17-O18
3	B	205	C8E	O9-C10-C11-O12
3	A	505	C8E	O9-C10-C11-O12
3	C	200	C8E	O9-C10-C11-O12
3	D	201	C8E	O18-C19-C20-O21
5	B	202	LBN	C26-C25-O5-C3
3	A	505	C8E	O18-C19-C20-O21
5	B	202	LBN	O6-C25-O5-C3
3	B	205	C8E	O18-C19-C20-O21
3	B	203	C8E	C4-C5-C6-C7
3	A	504	C8E	C3-C4-C5-C6
3	A	503	C8E	C2-C3-C4-C5
3	A	505	C8E	C2-C3-C4-C5
3	B	201	C8E	O18-C19-C20-O21
3	A	504	C8E	C2-C3-C4-C5
3	A	505	C8E	C1-C2-C3-C4
3	B	201	C8E	O12-C13-C14-O15
3	B	203	C8E	C3-C4-C5-C6
3	B	205	C8E	C16-C17-O18-C19
5	B	202	LBN	C1-C2-C3-O5
3	B	205	C8E	O15-C16-C17-O18
3	A	505	C8E	O12-C13-C14-O15
3	C	200	C8E	C3-C4-C5-C6
3	A	505	C8E	C4-C5-C6-C7
3	A	504	C8E	C4-C5-C6-C7

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	505	C8E	O15-C16-C17-O18
3	B	201	C8E	C17-C16-O15-C14
3	A	503	C8E	C1-C2-C3-C4
3	B	203	C8E	C6-C7-C8-O9
5	B	202	LBN	C26-C27-C28-C29
3	B	205	C8E	C1-C2-C3-C4
3	A	505	C8E	C3-C4-C5-C6
3	B	201	C8E	C1-C2-C3-C4
5	B	202	LBN	C31-C32-C33-C4
3	C	200	C8E	C17-C16-O15-C14
3	A	505	C8E	C5-C6-C7-C8
3	D	201	C8E	O9-C10-C11-O12
3	B	205	C8E	C3-C4-C5-C6
3	B	205	C8E	C14-C13-O12-C11
3	D	201	C8E	C16-C17-O18-C19
3	C	200	C8E	C11-C10-O9-C8
3	B	201	C8E	C10-C11-O12-C13
3	B	205	C8E	C13-C14-O15-C16
3	D	201	C8E	C17-C16-O15-C14
3	B	205	C8E	C20-C19-O18-C17
3	B	201	C8E	C2-C3-C4-C5
3	A	503	C8E	C3-C4-C5-C6
3	A	505	C8E	C6-C7-C8-O9
3	A	504	C8E	C7-C8-O9-C10
3	D	201	C8E	C7-C8-O9-C10
3	C	200	C8E	C10-C11-O12-C13
3	C	200	C8E	C7-C8-O9-C10
3	B	205	C8E	C10-C11-O12-C13
3	B	203	C8E	C2-C3-C4-C5
4	B	206	EDO	O1-C1-C2-O2
5	B	202	LBN	C35-C36-C37-C38
3	A	505	C8E	C16-C17-O18-C19
3	B	201	C8E	C20-C19-O18-C17
3	D	201	C8E	C13-C14-O15-C16
3	A	503	C8E	C4-C5-C6-C7
5	B	202	LBN	C9-O2-P1-O4
3	B	201	C8E	C7-C8-O9-C10
3	D	201	C8E	O12-C13-C14-O15


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







4.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

5 Fit of model and data [\(i\)](#)

5.1 Protein, DNA and RNA chains [\(i\)](#)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates [\(i\)](#)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands [\(i\)](#)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers [\(i\)](#)

EDS failed to run properly - this section is therefore empty.