



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 9, 2023 – 03:47 AM EDT

PDB ID : 6WPC
Title : Crystal structure of Bacillus thuringiensis Cry1A.2 tryptic core variant
Authors : Rydel, T.J.; Evdokimov, A.
Deposited on : 2020-04-27
Resolution : 2.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	591	
1	B	591	
1	C	591	
1	D	591	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 18743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cry1A.2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	588	4668	2970	816	874	8	0	1	0
1	B	588	4671	2972	817	874	8	0	1	0
1	C	588	4665	2968	816	873	8	0	0	0
1	D	588	4673	2973	819	873	8	0	1	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	16	Total	O	0	0
			16	16		
2	B	20	Total	O	0	0
			20	20		
2	C	13	Total	O	0	0
			13	13		
2	D	17	Total	O	0	0
			17	17		


3 Residue-property plots i


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cry1A.2





• Molecule 1: Cry1A.2






• Molecule 1: Cry1A.2



• Molecule 1: Cry1A.2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	120.58Å 225.82Å 239.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.12 – 2.99 46.12 – 2.99	Depositor EDS
% Data completeness (in resolution range)	94.5 (46.12-2.99) 94.8 (46.12-2.99)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 3.01Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660, PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.198 , 0.258 0.198 , 0.258	Depositor DCC
R_{free} test set	3189 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	45.8	Xtrriage
Anisotropy	0.057	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 21.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18743	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.48 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.3461e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	4/4786 (0.1%)	0.96	32/6509 (0.5%)
1	B	0.63	5/4789 (0.1%)	0.86	13/6513 (0.2%)
1	C	0.60	0/4780	0.82	10/6501 (0.2%)
1	D	0.61	1/4791 (0.0%)	0.94	28/6515 (0.4%)
All	All	0.62	10/19146 (0.1%)	0.90	83/26038 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	5
1	C	0	4
1	D	0	5
All	All	0	20

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	616	GLU	CG-CD	8.49	1.64	1.51
1	D	237	GLU	CB-CG	-6.22	1.40	1.52
1	A	332	ARG	CB-CG	-5.69	1.37	1.52
1	B	616	GLU	CD-OE2	5.52	1.31	1.25
1	A	557	ARG	CZ-NH2	-5.43	1.25	1.33

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	ARG	NE-CZ-NH2	-16.84	111.88	120.30
1	B	298	ASP	CB-CG-OD2	-15.64	104.22	118.30
1	D	332	ARG	NE-CZ-NH1	15.24	127.92	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	111	GLU	CA-CB-CG	-15.16	80.04	113.40
1	D	557	ARG	NE-CZ-NH2	-13.49	113.55	120.30

There are no chirality outliers.

5 of 20 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	126	ARG	Sidechain
1	A	149	GLU	Sidechain
1	A	392	PHE	Peptide
1	A	462	SER	Peptide
1	A	488	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4668	0	4547	97	1
1	B	4671	0	4552	106	0
1	C	4665	0	4544	105	0
1	D	4673	0	4557	107	1
2	A	16	0	0	2	0
2	B	20	0	0	2	0
2	C	13	0	0	2	0
2	D	17	0	0	3	0
All	All	18743	0	18200	411	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 411 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:616:GLU:OE1	1:B:630:LEU:HD23	1.09	1.24
1:B:616:GLU:OE1	1:B:630:LEU:CD2	2.02	1.07

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:552:SER:HB2	1:B:630:LEU:HD13	1.39	1.04
1:D:400:GLN:HB2	1:D:467:SER:HB2	1.44	0.99
1:B:547:ARG:NE	1:B:599:PHE:CZ	2.34	0.95

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:629:GLU:OE1	1:D:522:ARG:NH2[8_545]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	587/591 (99%)	549 (94%)	27 (5%)	11 (2%)	8	36
1	B	587/591 (99%)	551 (94%)	30 (5%)	6 (1%)	15	53
1	C	586/591 (99%)	542 (92%)	34 (6%)	10 (2%)	9	39
1	D	587/591 (99%)	548 (93%)	29 (5%)	10 (2%)	9	39
All	All	2347/2364 (99%)	2190 (93%)	120 (5%)	37 (2%)	9	40

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	393	ASN
1	A	564	ALA
1	A	601	ASN
1	B	396	ILE
1	C	171	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	505/508 (99%)	493 (98%)	12 (2%)	49 79
1	B	505/508 (99%)	492 (97%)	13 (3%)	46 78
1	C	504/508 (99%)	498 (99%)	6 (1%)	71 90
1	D	505/508 (99%)	486 (96%)	19 (4%)	33 69
All	All	2019/2032 (99%)	1969 (98%)	50 (2%)	49 79

5 of 50 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	345	SER
1	D	273	SER
1	D	624	SER
1	C	523	ARG
1	D	140	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	291	ASN
1	C	542	GLN
1	D	291	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	588/591 (99%)	-0.05	6 (1%) 82 59	19, 36, 65, 93	0
1	B	588/591 (99%)	-0.25	5 (0%) 84 63	19, 33, 58, 88	0
1	C	588/591 (99%)	-0.23	5 (0%) 84 63	20, 35, 57, 93	0
1	D	588/591 (99%)	-0.12	6 (1%) 82 59	19, 35, 63, 94	0
All	All	2352/2364 (99%)	-0.16	22 (0%) 84 63	19, 35, 61, 94	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	624	SER	8.1
1	B	624	SER	5.9
1	C	623	GLY	4.5
1	B	393	ASN	3.5
1	D	387	LEU	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.