

wwPDB X-ray Structure Validation Summary Report (i)

Dec 8, 2020 – 11:11 am GMT

PDB ID : 7ACG

Title : In meso structure of the alginate exporter, AlgE, from Pseudomonas aerugi-

nosa in 9.8 monoacylglycerol

Authors: Smithers, L.; van Dalsen, L.; Boland, C.; Caffrey, M.

Deposited on : 2020-09-10

Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.15.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.15.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			6%		
1	A	479	86%	10%	5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	${f Res}$	Chirality	Geometry	Clashes	Electron density
2	LH9	A	504	X	_	-	-





 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	LH9	A	506	X	-	-	-
2	LH9	A	513	X	-	-	-



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 7922 atoms, of which 3847 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Alginate biosynthesis protein AlgE.


Mol	Chain	Residues			Atom	S			ZeroOcc	AltConf	Trace
1	A	457	Total 7053	C 2279	H 3397	N 659	O 713	S 5	0	10	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	=	initiating methionine	UNP A0A080VSR9
A	-18	GLY	=	expression tag	UNP A0A080VSR9
A	-17	SER	-	expression tag	UNP A0A080VSR9
A	-16	SER	-	expression tag	UNP A0A080VSR9
A	-15	HIS	-	expression tag	UNP A0A080VSR9
A	-14	HIS	-	expression tag	UNP A0A080VSR9
A	-13	HIS	-	expression tag	UNP A0A080VSR9
A	-12	HIS	ı	expression tag	UNP A0A080VSR9
A	-11	HIS	-	expression tag	UNP A0A080VSR9
A	-10	HIS	-	expression tag	UNP A0A080VSR9
A	-9	SER	-	expression tag	UNP A0A080VSR9
A	-8	SER	-	expression tag	UNP A0A080VSR9
A	-7	GLY	-	expression tag	UNP A0A080VSR9
A	-6	LEU	-	expression tag	UNP A0A080VSR9
A	-5	VAL	-	expression tag	UNP A0A080VSR9
A	-4	PRO	=	expression tag	UNP A0A080VSR9
A	-3	ARG	-	expression tag	UNP A0A080VSR9
A	-2	GLY	=	expression tag	UNP A0A080VSR9
A	-1	SER	=	expression tag	UNP A0A080VSR9
A	0	HIS		expression tag	UNP A0A080VSR9
A	1	MET	-	expression tag	UNP A0A080VSR9

• Molecule 2 is $[(2 \{S\})-2,3-bis(oxidanyl)propyl]$ heptadec-9-enoate (three-letter code: LH9) (formula: $C_{20}H_{38}O_4$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	A	Aton	ns		ZeroOcc	AltConf
2	A	1	Total	С	Н	Ο	0	0
	A	1	62	20	38	4	0	0
2	A	1	Total	С	Η	О	0	0
	11	1	62	20	38	4	0	U
2	A	1	Total	\mathbf{C}	Η	Ο	0	0
	11	1	62	20	38	4		U
2	A	1	Total	\mathbf{C}	Η	Ο	0	0
	11	1	62	20	38	4		U
2	A	1	Total	\mathbf{C}	Η	Ο	0	0
	11	1	62	20	38	4	0	U
2	A	1	Total	\mathbf{C}	Η	Ο	0	0
	7.1	1	62	20	38	4		U
2	A	1	Total	\mathbf{C}	Η	Ο	0	0
	7.1	1	62	20	38	4		Ŭ
$\frac{1}{2}$	A	1	Total	С	Η	Ο	0	0
	71	1	62	20	38	4	0	Ů
2	A	1	Total	С	Η	Ο	0	0
	11	1	62	20	38	4		U
2	A	1	Total	\mathbf{C}	Η	Ο	0	0
	11	1	62	20	38	4		U


 \bullet Molecule 3 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $\mathrm{C_{10}H_{22}O_6}).$





Mol	Chain	Residues	A	Atoms				AltConf
9	Λ	1	Total	С	Н	О	0	0
3	A	1	37	10	21	6	0	0


 \bullet Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 5	O 4	S 1	0	0


• Molecule 5 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (three-letter code: C8E) (formula: $C_{16}H_{34}O_5$).





Mol	Chain	Residues	A	Atoms				AltConf
ĸ	Α	1	Total	С	Н	О	0	0
3	A	1	55	16	34	5	U	0


• Molecule 6 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



\mathbf{Mol}	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
6	A	1	Total 18	C 6	H 5	O 7	0	0

• Molecule 7 is 2-(2-METHOXYETHOXY)ETHANOL (three-letter code: PG0) (formula: $C_5H_{12}O_3$).





Mol	Chain	Residues	A	Atoms				AltConf
7	Α	1	Total	С	Н	О	0	0
1	A	1	18	5	10	3	U	0

 \bullet Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
8	A	2	Total Na 2 2	0	0

• Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	114	Total O 114 114	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alginate biosynthesis protein AlgE





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	56.75Å 74.19Å 114.99Å	Depositor
a, b, c, α , β , γ	90.00° 102.73° 90.00°	Depositor
Resolution (Å)	44.37 - 1.85	Depositor
resolution (A)	44.37 - 1.85	EDS
% Data completeness	99.1 (44.37-1.85)	Depositor
(in resolution range)	99.1 (44.37-1.85)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.67 (at 1.86Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
D D.	0.184 , 0.220	Depositor
R, R_{free}	0.183 , 0.220	DCC
R_{free} test set	1983 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtriage
Anisotropy	0.377	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.42, 50.5	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7922	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, 1PE, C8E, PG0, CIT, LH9, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
MIGI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.36	0/3802	0.63	0/5149	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3656	3397	3346	35	0
2	A	240	380	0	2	0
3	A	16	21	22	0	0
4	A	5	0	0	0	0
5	A	21	34	34	0	0
6	A	13	5	5	0	0
7	A	8	10	12	5	0
8	A	2	0	0	0	0
9	A	114	0	0	1	0
All	All	4075	3847	3419	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} & (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:316:THR:HG23	7:A:512:PG0:H51	1.47	0.97
1:A:427:ILE:HD13	1:A:455:PHE:CD1	2.24	0.72
1:A:337[B]:GLU:OE1	1:A:430:ARG:NH1	2.23	0.72
1:A:128:GLN:O	7:A:512:PG0:H53	1.90	0.71
1:A:315:GLN:HA	7:A:512:PG0:H52	1.76	0.66

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	${f Analysed}$	Favoured	Allowed	Outliers	Percentiles
1	A	459/479 (96%)	445 (97%)	14 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/388 (98%)	379 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are



no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Т	Chain	Dag	T : 1_	Во	Bond lengths			ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LH9	A	514	-	23,23,23	1.23	2 (8%)	24,24,24	0.93	1 (4%)
6	CIT	A	511	-	3,12,12	0.99	0	3,17,17	2.21	1 (33%)
2	LH9	A	513	-	23,23,23	1.09	2 (8%)	24,24,24	1.11	1 (4%)
7	PG0	A	512	-	7,7,7	0.51	0	6,6,6	0.37	0
2	LH9	A	517	_	23,23,23	1.11	2 (8%)	24,24,24	1.26	2 (8%)
4	SO4	A	508	-	4,4,4	0.14	0	6,6,6	0.21	0
2	LH9	A	504	_	23,23,23	1.29	2 (8%)	24,24,24	1.24	2 (8%)
2	LH9	A	505	-	23,23,23	1.11	2 (8%)	24,24,24	1.22	1 (4%)
2	LH9	A	506	-	23,23,23	1.13	2 (8%)	24,24,24	1.05	1 (4%)
2	LH9	A	501	-	23,23,23	1.13	3 (13%)	24,24,24	1.03	0
2	LH9	A	502	-	23,23,23	1.13	3 (13%)	24,24,24	1.23	1 (4%)
5	C8E	A	509	-	20,20,20	0.42	0	19,19,19	0.31	0
2	LH9	A	510	_	23,23,23	1.15	3 (13%)	24,24,24	0.83	0



7ACG

Mol	Tuno	Type Chain Res			Bond lengths			Bond angles		
MIOI	wor Type Chain Res L	Res Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2		
3	1PE	A	503	-	15,15,15	0.53	0	14,14,14	0.31	0
2	LH9	A	507	_	23,23,23	1.14	2 (8%)	24,24,24	0.97	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LH9	A	514	-	-	14/23/23/23	-
2	LH9	A	504	-	1/1/4/4	12/23/23/23	-
2	LH9	A	513	-	1/1/4/4	13/23/23/23	-
7	PG0	A	512	-	-	4/5/5/5	-
2	LH9	A	517	-	-	8/23/23/23	-
2	LH9	A	505	-	-	15/23/23/23	-
2	LH9	A	506	-	1/1/4/4	10/23/23/23	-
2	LH9	A	501	-	-	9/23/23/23	-
2	LH9	A	502	-	-	7/23/23/23	-
2	LH9	A	510	-	-	14/23/23/23	-
5	C8E	A	509	-	-	9/18/18/18	-
2	LH9	A	507	_	-	7/23/23/23	-
3	1PE	A	503		-	7/13/13/13	-
6	CIT	A	511	_	-	0/6/16/16	-

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	${ m Observed(\AA)}$	$\operatorname{Ideal}(ext{\AA})$
2	A	504	LH9	C9-C8	4.04	1.55	1.31
2	A	514	LH9	C9-C8	3.96	1.54	1.31
2	A	510	LH9	C9-C8	3.82	1.53	1.31
2	A	501	LH9	C9-C8	3.82	1.53	1.31
2	A	506	LH9	C9-C8	3.76	1.53	1.31

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
2	A	504	LH9	O2-C17-C16	3.56	123.08	111.91
2	A	517	LH9	O2-C17-C16	3.34	122.39	111.91
6	A	511	CIT	C3-C4-C5	-3.24	109.80	114.98

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	A	502	LH9	C15-C16-C17	-2.87	103.17	113.62
2	A	513	LH9	O2-C17-C16	2.87	120.92	111.91

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	513	LH9	C19
2	A	504	LH9	C19
2	A	506	LH9	C19

5 of 129 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	514	LH9	C18-C19-C20-O4
2	A	514	LH9	O3-C19-C20-O4
2	A	513	LH9	C18-C19-C20-O4
2	A	513	LH9	O3-C19-C20-O4
2	A	504	LH9	O2-C18-C19-C20


There are no ring outliers.

3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	512	PG0	5	0
2	A	504	LH9	1	0
2	A	501	LH9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	457/479 (95%)	0.21	28 (6%) 21 20	20, 32, 60, 104	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	213	LEU	6.6
1	A	86	ASP	5.4
1	A	210	VAL	4.8
1	A	212	ASN	4.4
1	A	265	ARG	4.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	${f Res}$	Atoms	RSCC	RSR	$\operatorname{\textbf{B-factors}}({ m \AA}^2)$	Q<0.9
2	LH9	A	514	24/24	0.59	0.28	55,72,85,87	0
2	LH9	A	505	24/24	0.63	0.27	39,61,76,82	0

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
2	LH9	A	504	24/24	0.64	0.24	48,69,83,99	0
2	LH9	A	506	24/24	0.74	0.26	35,60,79,94	0
2	LH9	A	502	24/24	0.77	0.22	34,57,78,83	0
2	LH9	A	517	24/24	0.80	0.17	37,58,73,83	0
2	LH9	A	510	24/24	0.80	0.37	$46,\!75,\!101,\!105$	0
2	LH9	A	507	24/24	0.80	0.27	40,58,76,90	0
2	LH9	A	501	24/24	0.81	0.23	35,48,100,114	0
2	LH9	A	513	24/24	0.84	0.28	49,75,97,99	0
5	C8E	A	509	21/21	0.84	0.17	49,68,82,93	0
3	1PE	A	503	16/16	0.88	0.21	32,56,85,86	0
7	PG0	A	512	8/8	0.90	0.11	$13,\!35,\!56,\!56$	0
8	NA	A	516	1/1	0.90	0.29	54,54,54,54	0
6	CIT	A	511	13/13	0.92	0.15	30,42,48,50	0
8	NA	A	515	1/1	0.98	0.05	30,30,30,30	0
4	SO4	A	508	5/5	0.99	0.06	27,28,35,35	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around LH9 A 514: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around LH9 A 505:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LH9 A 504:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LH9 A 506:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LH9 A 502:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LH9 A 517:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LH9 A 510:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LH9 A 507:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LH9 A 501:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around C8E A 509:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

