



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 5, 2021 – 12:15 PM GMT

PDB ID : 7B0H  
Title : TgoT\_6G12 Ternary complex  
Authors : Samson, C.; Legrand, P.; Tekpinar, M.; Rozenski, J.; Abramov, M.; Holliger, P.; Pinheiro, V.; Herdewijn, P.; Delarue, M.  
Deposited on : 2020-11-19  
Resolution : 3.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1665 (3.20-3.12)
Clashscore	141614	1804 (3.20-3.12)
Ramachandran outliers	138981	1770 (3.20-3.12)
Sidechain outliers	138945	1769 (3.20-3.12)
RSRZ outliers	127900	1616 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	13	15% (Poor fit) 69% (0-1 outliers) 31% (1-3 outliers)
1	C	13	31% (Poor fit) 69% (0-1 outliers) 31% (1-3 outliers)
2	D	6	83% (0-1 outliers) 17% (1-3 outliers)
2	G	6	67% (0-1 outliers) 33% (1-3 outliers)
3	E	773	7% (Poor fit) 84% (0-1 outliers) 8% (1-3 outliers) 8% (Not modelled)

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	773	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MG	F	803	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12977 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(P\*AP\*AP\*CP\*GP\*GP\*CP\*AP\*AP\*AP\*TP\*GP\*CP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	13	Total 267	C 127	N 56	O 72	P 12	0	0	0
1	C	13	Total 270	C 127	N 56	O 74	P 13	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(P\*CP\*GP\*CP\*AP\*TP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	D	6	Total 121	C 58	N 20	O 37	P 6	0	0	0
2	G	6	Total 121	C 58	N 20	O 37	P 6	0	0	0

- Molecule 3 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	713	Total 5870	C 3794	N 976	O 1086	S 14	0	0	0
3	F	762	Total 6264	C 4039	N 1062	O 1149	S 14	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	93	GLN	VAL	engineered mutation	UNP P56689
E	141	ALA	ASP	engineered mutation	UNP P56689
E	143	ALA	GLU	engineered mutation	UNP P56689
E	485	LEU	ALA	engineered mutation	UNP P56689
E	589	ALA	VAL	engineered mutation	UNP P56689
E	609	LYS	GLU	engineered mutation	UNP P56689
E	610	MET	ILE	engineered mutation	UNP P56689

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	659	GLN	LYS	engineered mutation	UNP P56689
E	664	GLN	GLU	engineered mutation	UNP P56689
E	665	PRO	GLN	engineered mutation	UNP P56689
E	668	LYS	ARG	engineered mutation	UNP P56689
E	669	GLN	ASP	engineered mutation	UNP P56689
E	671	HIS	LYS	engineered mutation	UNP P56689
E	674	ARG	LYS	engineered mutation	UNP P56689
E	676	ARG	THR	engineered mutation	UNP P56689
E	681	SER	ALA	engineered mutation	UNP P56689
E	704	PRO	LEU	engineered mutation	UNP P56689
E	730	GLY	GLU	engineered mutation	UNP P56689
F	93	GLN	VAL	engineered mutation	UNP P56689
F	141	ALA	ASP	engineered mutation	UNP P56689
F	143	ALA	GLU	engineered mutation	UNP P56689
F	485	LEU	ALA	engineered mutation	UNP P56689
F	589	ALA	VAL	engineered mutation	UNP P56689
F	609	LYS	GLU	engineered mutation	UNP P56689
F	610	MET	ILE	engineered mutation	UNP P56689
F	659	GLN	LYS	engineered mutation	UNP P56689
F	664	GLN	GLU	engineered mutation	UNP P56689
F	665	PRO	GLN	engineered mutation	UNP P56689
F	668	LYS	ARG	engineered mutation	UNP P56689
F	669	GLN	ASP	engineered mutation	UNP P56689
F	671	HIS	LYS	engineered mutation	UNP P56689
F	674	ARG	LYS	engineered mutation	UNP P56689
F	676	ARG	THR	engineered mutation	UNP P56689
F	681	SER	ALA	engineered mutation	UNP P56689
F	704	PRO	LEU	engineered mutation	UNP P56689
F	730	GLY	GLU	engineered mutation	UNP P56689


- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	F	2	Total Mn 2 2	0	0
4	E	2	Total Mn 2 2	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	F	1	Total Mg 1 1	0	0
5	E	1	Total Mg 1 1	0	0

- Molecule 6 is THYMIDINE-5'-TRIPHOSPHATE (three-letter code: TTP) (formula:  $C_{10}H_{17}N_2O_{14}P_3$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
6	E	1	Total	C	N	O	P	0	0
			29	10	2	14	3		
6	F	1	Total	C	N	O	P	0	0
			29	10	2	14	3		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(P\*AP\*AP\*CP\*GP\*GP\*CP\*AP\*AP\*AP\*TP\*GP\*CP\*G)-3')



- Molecule 1: DNA (5'-D(P\*AP\*AP\*CP\*GP\*GP\*CP\*AP\*AP\*AP\*TP\*GP\*CP\*G)-3')





- Molecule 2: DNA (5'-D(P\*CP\*GP\*CP\*AP\*TP\*T)-3')




- Molecule 2: DNA (5'-D(P\*CP\*GP\*CP\*AP\*TP\*T)-3')





- Molecule 3: DNA polymerase





● Molecule 3: DNA polymerase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.82Å 112.33Å 186.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.92 – 3.15 48.92 – 3.15	Depositor EDS
% Data completeness (in resolution range)	80.5 (48.92-3.15) 80.5 (48.92-3.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.52 (at 3.12Å)	Xtrriage
Refinement program	BUSTER 2.10.3 (20-MAY-2020)	Depositor
R, $R_{free}$	0.232 , 0.247 0.256 , 0.266	Depositor DCC
$R_{free}$ test set	1648 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	88.7	Xtrriage
Anisotropy	0.068	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 50.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.033 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	12977	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	104.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MN, TTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/301	0.86	0/463
1	C	0.71	0/304	0.79	0/467
2	D	0.64	0/134	0.82	0/204
2	G	0.68	0/134	0.87	0/204
3	E	0.27	0/6004	0.50	0/8098
3	F	0.27	0/6408	0.51	2/8641 (0.0%)
All	All	0.32	0/13285	0.53	2/18077 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	665	PRO	N-CA-CB	-5.46	96.60	102.60
3	F	665	PRO	CA-N-CD	-5.45	103.88	111.50

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	267	0	146	7	0
1	C	270	0	145	3	0
2	D	121	0	69	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	G	121	0	69	3	0
3	E	5870	0	5883	39	0
3	F	6264	0	6304	67	0
4	E	2	0	0	0	0
4	F	2	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
6	E	29	0	13	0	0
6	F	29	0	13	0	0
All	All	12977	0	12642	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:676:ARG:NE	3:F:684:LYS:NZ	2.02	1.06
3:F:676:ARG:NE	3:F:684:LYS:HZ2	1.62	0.94
3:F:689:ARG:NH2	3:F:691:ILE:HD11	1.87	0.90
3:F:683:ALA:O	3:F:693:ILE:CD1	2.24	0.85
3:F:676:ARG:CD	3:F:684:LYS:HZ2	1.93	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	707/773 (92%)	685 (97%)	22 (3%)	0	100	100
3	F	760/773 (98%)	735 (97%)	25 (3%)	0	100	100
All	All	1467/1546 (95%)	1420 (97%)	47 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	E	621/668 (93%)	613 (99%)	8 (1%)	69	86
3	F	660/668 (99%)	655 (99%)	5 (1%)	81	92
All	All	1281/1336 (96%)	1268 (99%)	13 (1%)	76	89

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	E	663	TYR
3	E	736	GLN
3	F	494	TYR
3	E	613	ARG
3	F	466	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	F	436	GLN
3	F	491	ASN
3	F	664	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	13/13 (100%)	0.99	2 (15%) 2 1	60, 81, 153, 157	0
1	C	13/13 (100%)	1.87	4 (30%) 0 0	86, 108, 204, 208	0
2	D	6/6 (100%)	0.07	0 100 100	95, 102, 124, 131	0
2	G	6/6 (100%)	0.67	0 100 100	87, 97, 115, 119	0
3	E	713/773 (92%)	0.51	53 (7%) 14 8	50, 103, 166, 184	0
3	F	762/773 (98%)	0.38	27 (3%) 44 27	70, 100, 129, 171	0
All	All	1513/1584 (95%)	0.46	86 (5%) 23 12	50, 101, 153, 208	0

The worst 5 of 86 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	673	TYR	9.4
3	F	670	LEU	8.8
3	E	295	ILE	8.1
1	C	-2	DA	7.9
3	E	305	LEU	6.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MN	E	801	1/1	0.75	0.13	165,165,165,165	0
5	MG	F	803	1/1	0.78	0.41	96,96,96,96	0
5	MG	E	803	1/1	0.89	0.26	80,80,80,80	0
6	TTP	E	804	29/29	0.92	0.24	93,94,96,96	0
4	MN	F	801	1/1	0.94	0.26	118,118,118,118	0
4	MN	E	802	1/1	0.95	0.09	100,100,100,100	0
6	TTP	F	804	29/29	0.95	0.28	67,69,69,69	0
4	MN	F	802	1/1	0.98	0.16	71,71,71,71	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








**Electron density around MG F 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MG E 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around TTP E 804:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MN F 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MN E 802:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around TTP F 804:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.