



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 7, 2023 – 05:01 PM EST

PDB ID : 7FSG
Title : SDCBP PanDDA analysis group deposition – The PDZ domains of SDCBP in complex with Z56772132
Authors : Bradshaw, W.J.; Katis, V.L.; Bountra, C.; von Delft, F.; Brennan, P.E.
Deposited on : 2023-01-24
Resolution : 2.02 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.32.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.1

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6478 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Syntenin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	191	1481	935	262	275	9	0	1	0
1	B	193	1495	943	264	279	9	0	1	0
1	C	193	1514	953	270	282	9	0	3	0
1	D	191	1498	944	265	280	9	0	3	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	104	SER	-	expression tag	UNP O00560
A	105	MET	-	expression tag	UNP O00560
B	104	SER	-	expression tag	UNP O00560
B	105	MET	-	expression tag	UNP O00560
C	104	SER	-	expression tag	UNP O00560
C	105	MET	-	expression tag	UNP O00560
D	104	SER	-	expression tag	UNP O00560
D	105	MET	-	expression tag	UNP O00560

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is D-GLUTAMIC ACID (three-letter code: DGL) (formula: C₅H₉NO₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			10	5	1	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	D	1	10	5	1	4	0	0

- Molecule 5 is GLYCINE (three-letter code: GLY) (formula: C₂H₅NO₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	C	1	5	2	1	2	0	0
5	D	1	5	2	1	2	0	0
5	D	1	5	2	1	2	0	0
5	D	1	5	2	1	2	0	0

- Molecule 6 is (5R)-5-[2-(4-methoxyphenyl)ethyl]-5-methylimidazolidine-2,4-dione (three-letter code: YG9) (formula: C₁₃H₁₆N₂O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	D	1	18	13	2	3	0	0
6	D	1	18	13	2	3	0	0

- Molecule 7 is ALANINE (three-letter code: ALA) (formula: $C_3H_7NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
7	D	1	6	3	1	2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	44	Total 44	O 44	0	0
8	B	113	Total 113	O 113	0	1
8	C	90	Total 91	O 91	0	1
8	D	113	Total 113	O 113	0	0


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Syntenin-1




- Molecule 1: Syntenin-1



- Molecule 1: Syntenin-1



- Molecule 1: Syntenin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.04Å 49.51Å 115.58Å 90.00° 94.74° 90.00°	Depositor
Resolution (Å)	115.18 – 2.02 115.18 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.7 (115.18-2.02) 99.8 (115.18-2.02)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 2.02Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.201 , 0.252 0.213 , 0.263	Depositor DCC
R_{free} test set	2936 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	36.2	Xtrriage
Anisotropy	0.184	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 38.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6478	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.23 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2988e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: YG9, DGL, EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/1502	0.98	2/2019 (0.1%)
1	B	0.84	0/1516	1.00	1/2038 (0.0%)
1	C	0.82	0/1535	0.98	1/2063 (0.0%)
1	D	0.91	2/1519 (0.1%)	0.99	2/2042 (0.1%)
All	All	0.85	2/6072 (0.0%)	0.99	6/8162 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	240	GLU	C-O	5.23	1.33	1.23
1	D	192	ASP	C-O	5.07	1.32	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	197	ARG	NE-CZ-NH2	-7.43	116.58	120.30
1	D	153	ARG	NE-CZ-NH1	-6.90	116.85	120.30
1	A	266	THR	CA-CB-OG1	-5.87	96.68	109.00
1	D	153	ARG	CG-CD-NE	-5.54	100.16	111.80
1	C	153	ARG	CG-CD-NE	-5.42	100.41	111.80
1	B	197	ARG	NE-CZ-NH2	-5.36	117.62	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	111	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1481	0	1530	12	0
1	B	1495	0	1541	5	0
1	C	1514	0	1558	17	1
1	D	1498	0	1540	12	0
2	A	4	0	6	0	0
2	B	8	0	12	0	0
2	C	12	0	18	0	0
2	D	8	0	12	0	0
3	B	5	0	0	0	0
3	D	10	0	0	0	0
4	B	10	0	7	0	0
4	D	10	0	7	0	0
5	C	5	0	2	0	0
5	D	15	0	6	0	0
6	D	36	0	0	0	0
7	D	6	0	4	0	0
8	A	44	0	0	1	0
8	B	113	0	0	2	0
8	C	91	0	0	3	0
8	D	113	0	0	2	0
All	All	6478	0	6243	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:108:ILE:HD12	1:C:198:THR:HG23	1.65	0.76
1:D:117:LEU:HD12	1:D:117:LEU:H	1.54	0.72
1:D:256[B]:ASP:O	1:D:260:THR:HG23	1.87	0.71
1:C:206:THR:OG1	1:C:208:HIS:ND1	2.29	0.66
1:B:202:HIS:O	8:B:401:HOH:O	2.15	0.64
1:C:206:THR:HG1	1:C:208:HIS:CE1	2.18	0.62
1:D:117:LEU:HD22	1:D:125:ILE:HG23	1.80	0.61
1:A:109:LYS:O	1:A:111:GLY:N	2.33	0.60
1:B:150:VAL:O	8:B:402:HOH:O	2.17	0.59
1:D:215:ASN:O	8:D:601:HOH:O	2.17	0.59
1:A:151:GLY:O	1:A:153:ARG:NH1	2.37	0.57
1:A:140:LEU:HD11	1:A:282:MET:SD	2.45	0.56
1:A:108:ILE:N	1:A:108:ILE:HD13	2.21	0.55
1:D:117:LEU:HD12	1:D:117:LEU:N	2.21	0.55
1:D:210:GLY:HA3	1:D:225:SER:HB2	1.89	0.55
1:D:286:ILE:HG23	1:D:290:LEU:HD12	1.89	0.54
1:D:210:GLY:HA3	1:D:225:SER:CB	2.38	0.53
1:C:182:PHE:O	1:C:183:GLY:O	2.29	0.51
1:C:110:GLN:HG3	1:C:111:GLY:H	1.76	0.50
1:B:294:THR:HG22	1:B:295:ILE:O	2.11	0.50
1:D:154:PHE:CD1	1:D:247:ILE:CD1	2.95	0.49
1:A:140:LEU:CD1	1:A:282:MET:SD	3.01	0.49
1:A:201:MET:CE	1:A:267:ILE:HD12	2.43	0.48
1:C:176:LYS:HE2	1:C:180:GLN:NE2	2.28	0.48
1:B:125:ILE:HD11	1:B:186:ILE:CD1	2.44	0.47
1:C:210:GLY:HA3	1:C:225:SER:CB	2.44	0.47
1:C:229:ARG:NH2	8:C:408:HOH:O	2.47	0.47
1:C:210:GLY:HA3	1:C:225:SER:HB2	1.95	0.47
1:A:244:GLN:HE21	1:A:257:ILE:HD13	1.79	0.46
1:A:223:LYS:CE	8:A:409:HOH:O	2.63	0.46
1:D:142:GLN:HB3	1:D:145:SER:HB3	1.98	0.45
1:D:197:ARG:NH2	8:D:602:HOH:O	2.34	0.45
1:C:273[B]:PHE:CD1	8:C:423:HOH:O	2.70	0.45
1:A:203:LYS:NZ	1:A:261:SER:O	2.41	0.44
1:C:121:GLN:HG2	8:C:456:HOH:O	2.17	0.44
1:C:240:GLU:HA	1:C:244:GLN:O	2.19	0.42
1:A:291:MET:O	1:A:293:HIS:CE1	2.73	0.42
1:A:235:GLU:OE1	1:A:235:GLU:HA	2.19	0.42
1:C:291:MET:HB2	1:C:291:MET:HE2	1.97	0.42
1:A:210:GLY:HA3	1:A:225:SER:HB2	2.02	0.42
1:C:157:GLN:HG2	1:C:159:LEU:HD12	2.02	0.42
1:C:160:GLN:HA	1:C:164:GLU:O	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:ILE:HD11	1:B:186:ILE:HD13	2.01	0.41
1:D:198:THR:HA	1:D:267:ILE:O	2.20	0.41
1:C:197[A]:ARG:HH11	1:C:197[A]:ARG:HD2	1.74	0.41
1:C:182:PHE:O	1:C:183:GLY:C	2.59	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:128:ARG:NH2	1:C:297:GLU:OE1[2_544]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/195 (97%)	180 (95%)	8 (4%)	2 (1%)	14	7
1	B	192/195 (98%)	188 (98%)	4 (2%)	0	100	100
1	C	194/195 (100%)	186 (96%)	7 (4%)	1 (0%)	29	22
1	D	192/195 (98%)	188 (98%)	3 (2%)	1 (0%)	29	22
All	All	768/780 (98%)	742 (97%)	22 (3%)	4 (0%)	29	22

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	110	GLN
1	A	183	GLY
1	C	183	GLY
1	D	262	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/168 (99%)	163 (98%)	3 (2%)	59	61
1	B	167/168 (99%)	163 (98%)	4 (2%)	49	49
1	C	169/168 (101%)	168 (99%)	1 (1%)	86	89
1	D	168/168 (100%)	164 (98%)	4 (2%)	49	49
All	All	670/672 (100%)	658 (98%)	12 (2%)	59	61

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	205	SER
1	A	223	LYS
1	A	259	SER
1	B	110	GLN
1	B	114	GLU
1	B	140	LEU
1	B	284	PRO
1	C	209	VAL
1	D	117	LEU
1	D	121	GLN
1	D	139	GLN
1	D	297	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	110	GLN
1	A	215	ASN
1	B	142	GLN
1	C	110	GLN
1	D	215	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	B	301	-	3,3,3	0.26	0	2,2,2	0.30	0
7	ALA	D	507	-	5,5,5	0.83	0	6,6,6	1.49	1 (16%)
4	DGL	D	508	-	8,9,9	0.97	0	10,11,11	0.99	0
3	SO4	D	506	-	4,4,4	0.33	0	6,6,6	0.10	0
6	YG9	D	501	-	17,19,19	2.57	5 (29%)	21,27,27	2.59	3 (14%)
4	DGL	B	304	-	8,9,9	1.08	1 (12%)	10,11,11	1.74	2 (20%)
2	EDO	C	302	-	3,3,3	0.38	0	2,2,2	0.44	0
2	EDO	D	504	-	3,3,3	0.37	0	2,2,2	0.13	0
2	EDO	C	301	-	3,3,3	0.25	0	2,2,2	0.35	0
2	EDO	B	302	-	3,3,3	0.22	0	2,2,2	0.28	0
5	GLY	D	511	-	4,4,4	0.97	0	3,4,4	1.67	1 (33%)
6	YG9	D	502	-	17,19,19	2.40	4 (23%)	21,27,27	2.60	3 (14%)
5	GLY	C	304	-	4,4,4	0.82	0	3,4,4	1.53	0
2	EDO	C	303	-	3,3,3	0.43	0	2,2,2	0.66	0
5	GLY	D	509	-	4,4,4	1.25	1 (25%)	3,4,4	1.35	0
2	EDO	D	503	-	3,3,3	0.28	0	2,2,2	0.41	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GLY	D	510	-	4,4,4	0.98	0	3,4,4	1.56	1 (33%)
3	SO4	D	505	-	4,4,4	0.35	0	6,6,6	0.06	0
3	SO4	B	303	-	4,4,4	0.36	0	6,6,6	0.04	0
2	EDO	A	301	-	3,3,3	0.18	0	2,2,2	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	301	-	-	0/1/1/1	-
2	EDO	D	504	-	-	1/1/1/1	-
7	ALA	D	507	-	-	4/4/4/4	-
5	GLY	C	304	-	-	0/2/2/2	-
2	EDO	C	303	-	-	1/1/1/1	-
2	EDO	C	301	-	-	0/1/1/1	-
5	GLY	D	509	-	-	2/2/2/2	-
2	EDO	D	503	-	-	0/1/1/1	-
4	DGL	D	508	-	-	4/9/9/9	-
2	EDO	B	302	-	-	0/1/1/1	-
5	GLY	D	510	-	-	2/2/2/2	-
6	YG9	D	502	-	-	1/8/23/23	0/2/2/2
6	YG9	D	501	-	-	2/8/23/23	0/2/2/2
4	DGL	B	304	-	-	1/9/9/9	-
5	GLY	D	511	-	-	0/2/2/2	-
2	EDO	C	302	-	-	0/1/1/1	-
2	EDO	A	301	-	-	0/1/1/1	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	502	YG9	C12-N11	7.45	1.48	1.35
6	D	501	YG9	C12-N11	6.55	1.47	1.35
6	D	501	YG9	C15-N14	5.44	1.45	1.37
6	D	501	YG9	C12-N14	4.82	1.48	1.39
6	D	502	YG9	C15-N14	3.86	1.42	1.37
6	D	502	YG9	C12-N14	3.70	1.46	1.39
6	D	502	YG9	O16-C15	-2.44	1.18	1.22
5	D	509	GLY	OXT-C	-2.38	1.22	1.30
4	B	304	DGL	OXT-C	-2.16	1.23	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	501	YG9	O16-C15	-2.12	1.19	1.22
6	D	501	YG9	C10-C09	2.04	1.57	1.53

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	502	YG9	C15-C09-N11	10.01	107.33	100.30
6	D	501	YG9	C15-C09-N11	9.88	107.24	100.30
6	D	502	YG9	C09-N11-C12	-5.34	107.77	112.77
4	B	304	DGL	OXT-C-O	-3.77	115.53	124.09
6	D	501	YG9	C09-N11-C12	-3.75	109.26	112.77
6	D	501	YG9	C15-N14-C12	-3.43	108.10	111.69
7	D	507	ALA	OXT-C-CA	2.62	123.51	114.06
4	B	304	DGL	OXT-C-CA	2.48	121.84	113.38
6	D	502	YG9	C15-N14-C12	-2.22	109.38	111.69
5	D	510	GLY	OXT-C-CA	2.13	121.91	113.45
5	D	511	GLY	OXT-C-O	-2.05	118.18	123.30

There are no chirality outliers.


All (18) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	D	502	YG9	C07-C08-C09-C15
6	D	501	YG9	C04-C03-O02-C01
6	D	501	YG9	C18-C03-O02-C01
5	D	509	GLY	OXT-C-CA-N
5	D	510	GLY	OXT-C-CA-N
5	D	510	GLY	O-C-CA-N
5	D	509	GLY	O-C-CA-N
7	D	507	ALA	O-C-CA-CB
7	D	507	ALA	OXT-C-CA-CB
2	C	303	EDO	O1-C1-C2-O2
4	D	508	DGL	OXT-C-CA-CB
4	D	508	DGL	O-C-CA-CB
7	D	507	ALA	OXT-C-CA-N
4	D	508	DGL	CA-CB-CG-CD
2	D	504	EDO	O1-C1-C2-O2
7	D	507	ALA	O-C-CA-N
4	B	304	DGL	N-CA-CB-CG
4	D	508	DGL	N-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/195 (97%)	0.25	20 (10%) 6 5	32, 47, 80, 99	20 (10%)
1	B	193/195 (98%)	-0.16	4 (2%) 63 63	27, 40, 58, 71	5 (2%)
1	C	193/195 (98%)	0.17	24 (12%) 4 3	26, 42, 67, 89	21 (10%)
1	D	191/195 (97%)	-0.08	8 (4%) 36 35	26, 36, 56, 80	14 (7%)
All	All	768/780 (98%)	0.04	56 (7%) 15 14	26, 41, 68, 99	60 (7%)

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	125	ILE	6.9
1	A	108	ILE	5.7
1	A	182	PHE	5.4
1	A	281	ARG	4.6
1	C	298	VAL	4.5
1	A	110	GLN	4.2
1	C	150	VAL	4.1
1	C	261	SER	4.0
1	C	211	PHE	3.9
1	A	112	ILE	3.9
1	D	125	ILE	3.9
1	A	111	GLY	3.8
1	C	197[A]	ARG	3.7
1	A	282	MET	3.5
1	C	109	LYS	3.4
1	C	294	THR	3.4
1	A	283	ALA	3.4
1	C	110	GLN	3.4
1	B	125	ILE	3.3
1	C	181	ALA	3.3
1	D	118	CYS	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	212	ILE	3.2
1	A	109	LYS	3.1
1	A	184	GLU	3.1
1	C	206	THR	3.1
1	D	184	GLU	3.1
1	C	149	LEU	3.1
1	A	286	ILE	3.0
1	C	182	PHE	2.9
1	C	106	ALA	2.9
1	D	261	SER	2.9
1	C	121	GLN	2.8
1	A	298	VAL	2.8
1	B	109	LYS	2.8
1	A	287	MET	2.8
1	C	297	GLU	2.8
1	A	258	LEU	2.7
1	D	160[A]	GLN	2.7
1	B	209	VAL	2.7
1	C	108	ILE	2.6
1	A	213	PHE	2.5
1	B	273[A]	PHE	2.5
1	C	209	VAL	2.5
1	C	107	GLU	2.5
1	C	118	CYS	2.5
1	A	290	LEU	2.4
1	D	117	LEU	2.3
1	A	212	ILE	2.3
1	A	215	ASN	2.2
1	D	108	ILE	2.2
1	C	184	GLU	2.2
1	C	183	GLY	2.2
1	A	262	GLY	2.1
1	C	185	LYS	2.1
1	D	124	LYS	2.1
1	A	263	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	C	303	4/4	0.74	0.18	58,63,69,70	0
7	ALA	D	507	6/6	0.78	0.35	56,71,77,78	0
5	GLY	D	511	5/5	0.81	0.17	60,61,64,66	0
4	DGL	D	508	10/10	0.81	0.16	43,68,74,76	0
5	GLY	D	510	5/5	0.82	0.28	48,58,59,63	0
6	YG9	D	502	18/18	0.87	0.16	27,28,30,30	18
2	EDO	C	302	4/4	0.87	0.12	39,40,43,46	0
3	SO4	B	303	5/5	0.90	0.18	95,95,106,107	0
3	SO4	D	505	5/5	0.92	0.20	71,81,88,91	0
6	YG9	D	501	18/18	0.93	0.15	26,28,30,30	18
2	EDO	D	503	4/4	0.94	0.12	26,26,26,30	0
5	GLY	C	304	5/5	0.94	0.11	47,48,50,54	0
5	GLY	D	509	5/5	0.94	0.13	68,69,71,75	0
4	DGL	B	304	10/10	0.94	0.13	44,61,67,68	0
2	EDO	D	504	4/4	0.95	0.13	35,37,37,37	0
2	EDO	C	301	4/4	0.95	0.13	34,36,38,40	0
2	EDO	A	301	4/4	0.95	0.11	45,46,47,49	0
3	SO4	D	506	5/5	0.96	0.19	57,59,76,78	0
2	EDO	B	302	4/4	0.97	0.10	32,38,42,43	0
2	EDO	B	301	4/4	0.97	0.13	34,38,39,42	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around YG9 D 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around YG9 D 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.