



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 10, 2023 – 08:04 AM EDT

PDB ID : 7T1V  
Title : Crystal structure of an equine H7 hemagglutinin from A/equine/NY/49/73 (H7N7) in complex with 3'-GcLN  
Authors : Zhu, X.; Wilson, I.A.  
Deposited on : 2021-12-02  
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	
1	C	334	
1	E	334	
2	B	186	
2	D	186	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	186	 77% 79% 11% • 9%
3	G	3	 67% 33%
4	H	3	 100%
5	I	2	 50% 50%
5	K	2	 50% 50%
6	J	2	 100%
6	L	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	NAG	E	401	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 12574 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	315	2448	1531	435	467	15	0	0	0
1	C	315	2448	1531	435	467	15	0	0	0
1	E	315	2448	1531	435	467	15	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	ASP	-	expression tag	UNP A0A348FV55
A	9	PRO	-	expression tag	UNP A0A348FV55
A	10	GLY	-	expression tag	UNP A0A348FV55
C	8	ASP	-	expression tag	UNP A0A348FV55
C	9	PRO	-	expression tag	UNP A0A348FV55
C	10	GLY	-	expression tag	UNP A0A348FV55
E	8	ASP	-	expression tag	UNP A0A348FV55
E	9	PRO	-	expression tag	UNP A0A348FV55
E	10	GLY	-	expression tag	UNP A0A348FV55

- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	170	1390	862	247	276	5	0	0	0
2	D	170	1390	862	247	276	5	0	0	0
2	F	170	1390	862	247	276	5	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	179	GLU	-	expression tag	UNP A0A348FV55
B	180	SER	-	expression tag	UNP A0A348FV55
B	181	GLY	-	expression tag	UNP A0A348FV55
B	182	ARG	-	expression tag	UNP A0A348FV55
B	183	LEU	-	expression tag	UNP A0A348FV55
B	184	VAL	-	expression tag	UNP A0A348FV55
B	185	PRO	-	expression tag	UNP A0A348FV55
B	186	ARG	-	expression tag	UNP A0A348FV55
D	179	GLU	-	expression tag	UNP A0A348FV55
D	180	SER	-	expression tag	UNP A0A348FV55
D	181	GLY	-	expression tag	UNP A0A348FV55
D	182	ARG	-	expression tag	UNP A0A348FV55
D	183	LEU	-	expression tag	UNP A0A348FV55
D	184	VAL	-	expression tag	UNP A0A348FV55
D	185	PRO	-	expression tag	UNP A0A348FV55
D	186	ARG	-	expression tag	UNP A0A348FV55
F	179	GLU	-	expression tag	UNP A0A348FV55
F	180	SER	-	expression tag	UNP A0A348FV55
F	181	GLY	-	expression tag	UNP A0A348FV55
F	182	ARG	-	expression tag	UNP A0A348FV55
F	183	LEU	-	expression tag	UNP A0A348FV55
F	184	VAL	-	expression tag	UNP A0A348FV55
F	185	PRO	-	expression tag	UNP A0A348FV55
F	186	ARG	-	expression tag	UNP A0A348FV55

- Molecule 3 is an oligosaccharide called N-glycolyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
3	G	3	47	25	2	20	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	H	3	39	22	2	15	0	0	0

- Molecule 5 is an oligosaccharide called N-glycolyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	I	2	33	17	1	15	0	0	0
5	K	2	33	17	1	15	0	0	0

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	J	2	28	16	2	10	0	0	0
6	L	2	28	16	2	10	0	0	0

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
7	A	1	14	8	1	5	0	0
7	C	1	14	8	1	5	0	0
7	E	1	14	8	1	5	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
8	A	298	298	298	0	0
8	B	87	87	87	0	0
8	C	267	267	267	0	0
8	D	84	84	84	0	0
8	E	62	62	62	0	0
8	F	12	12	12	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain



- Molecule 2: Hemagglutinin HA2 chain











PRO  
ARG

- Molecule 2: Hemagglutinin HA2 chain



- Molecule 2: Hemagglutinin HA2 chain



- Molecule 3: N-glycolyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1  
GAL2  
MGCC3

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1  
MAG2  
BMA3

- Molecule 5: N-glycolyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose



GAL1  
MGC2


- Molecule 5: N-glycolyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose

Chain K:  50% 50%GAL1  
MGC2

- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  100%MAG1  
MAG2

- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  100%MAG1  
MAG2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.75Å 112.75Å 130.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.01 – 2.05 48.82 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.01-2.05) 99.3 (48.82-2.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.217 , 0.245 0.221 , 0.246	Depositor DCC
$R_{free}$ test set	5856 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.7	Xtriage
Anisotropy	0.440	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 53.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.010 for -h,-k,l 0.027 for h,-h-k,-l 0.019 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12574	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, BMA, NGC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/2497	0.73	0/3368
1	C	0.48	0/2497	0.72	1/3368 (0.0%)
1	E	0.47	0/2497	0.84	8/3368 (0.2%)
2	B	0.52	0/1415	0.72	3/1907 (0.2%)
2	D	0.45	0/1415	0.64	1/1907 (0.1%)
2	F	0.43	0/1415	0.64	2/1907 (0.1%)
All	All	0.48	0/11736	0.73	15/15825 (0.1%)

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	193	ARG	NE-CZ-NH1	9.20	124.90	120.30
1	E	155	LEU	CB-CG-CD1	-8.18	97.10	111.00
2	F	123	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	E	52	CYS	CB-CA-C	6.20	122.80	110.40
2	D	123	ARG	NE-CZ-NH1	6.15	123.38	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2448	0	2419	4	0
1	C	2448	0	2419	11	0
1	E	2448	0	2419	28	0
2	B	1390	0	1303	6	0
2	D	1390	0	1303	9	0
2	F	1390	0	1303	11	0
3	G	47	0	31	0	0
4	H	39	0	34	0	0
5	I	33	0	19	1	0
5	K	33	0	19	0	0
6	J	28	0	25	0	0
6	L	28	0	25	0	0
7	A	14	0	13	0	0
7	C	14	0	13	0	0
7	E	14	0	13	0	0
8	A	298	0	0	1	0
8	B	87	0	0	2	0
8	C	267	0	0	1	0
8	D	84	0	0	2	0
8	E	62	0	0	0	0
8	F	12	0	0	0	0
All	All	12574	0	11358	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 65 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:81:GLU:OE2	1:C:119:LYS:HD2	1.89	0.72
1:E:132:THR:HG23	1:E:152:LYS:HD2	1.70	0.71
2:B:30:GLN:HE22	2:B:145:ASP:HB2	1.56	0.70
1:A:150:GLU:OE1	1:A:256:ARG:HD3	1.94	0.66
1:C:150:GLU:OE1	1:C:256:ARG:HD3	1.95	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/334 (94%)	306 (98%)	7 (2%)	0	100	100
1	C	313/334 (94%)	306 (98%)	7 (2%)	0	100	100
1	E	313/334 (94%)	307 (98%)	6 (2%)	0	100	100
2	B	168/186 (90%)	157 (94%)	11 (6%)	0	100	100
2	D	168/186 (90%)	157 (94%)	11 (6%)	0	100	100
2	F	168/186 (90%)	156 (93%)	12 (7%)	0	100	100
All	All	1443/1560 (92%)	1389 (96%)	54 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/288 (94%)	262 (97%)	8 (3%)	41	34
1	C	270/288 (94%)	262 (97%)	8 (3%)	41	34
1	E	270/288 (94%)	256 (95%)	14 (5%)	23	14
2	B	146/160 (91%)	141 (97%)	5 (3%)	37	30
2	D	146/160 (91%)	142 (97%)	4 (3%)	44	38
2	F	146/160 (91%)	141 (97%)	5 (3%)	37	30
All	All	1248/1344 (93%)	1204 (96%)	44 (4%)	36	29

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	44	GLU
1	E	277	CYS
1	E	132	THR
1	E	172	LYS
1	E	310	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	163	GLN
2	B	159	HIS
1	C	163	GLN
2	F	62	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	G	1	3	15,15,15	0.55	0	21,21,21	0.88	0
3	GAL	G	2	3	11,11,12	0.27	0	15,15,17	0.86	0
3	NGC	G	3	3	21,21,22	0.91	1 (4%)	25,29,32	1.24	1 (4%)
4	NAG	H	1	4,2	14,14,15	0.39	0	17,19,21	0.93	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	H	2	4	14,14,15	0.32	0	17,19,21	0.56	0
4	BMA	H	3	4	11,11,12	0.39	0	15,15,17	0.89	0
5	GAL	I	1	5	12,12,12	0.68	0	17,17,17	1.21	2 (11%)
5	NGC	I	2	5	21,21,22	0.93	1 (4%)	25,29,32	1.19	2 (8%)
6	NAG	J	1	2,6	14,14,15	0.33	0	17,19,21	0.95	0
6	NAG	J	2	6	14,14,15	0.35	0	17,19,21	0.81	0
5	GAL	K	1	5	12,12,12	0.49	0	17,17,17	0.61	0
5	NGC	K	2	5	21,21,22	0.60	0	25,29,32	1.13	3 (12%)
6	NAG	L	1	2,6	14,14,15	0.40	0	17,19,21	1.23	2 (11%)
6	NAG	L	2	6	14,14,15	0.37	0	17,19,21	1.24	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3	-	0/6/26/26	0/1/1/1
3	GAL	G	2	3	-	1/2/19/22	0/1/1/1
3	NGC	G	3	3	-	1/19/36/40	0/1/1/1
4	NAG	H	1	4,2	-	1/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1
4	BMA	H	3	4	-	2/2/19/22	0/1/1/1
5	GAL	I	1	5	-	1/2/22/22	0/1/1/1
5	NGC	I	2	5	-	1/19/36/40	0/1/1/1
6	NAG	J	1	2,6	-	2/6/23/26	0/1/1/1
6	NAG	J	2	6	-	0/6/23/26	0/1/1/1
5	GAL	K	1	5	-	0/2/22/22	0/1/1/1
5	NGC	K	2	5	-	1/19/36/40	0/1/1/1
6	NAG	L	1	2,6	-	2/6/23/26	0/1/1/1
6	NAG	L	2	6	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	I	2	NGC	C2-C1	3.00	1.55	1.52
3	G	3	NGC	O6-C2	-2.30	1.40	1.43

The worst 5 of 12 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	L	1	NAG	C1-O5-C5	3.54	116.99	112.19
3	G	3	NGC	C6-O6-C2	3.41	118.64	111.34
6	L	2	NAG	C1-C2-N2	-3.06	105.26	110.49
5	K	2	NGC	C4-C5-N5	-2.82	104.79	110.38
5	I	2	NGC	O1B-C1-C2	2.71	120.77	113.03

There are no chirality outliers.

5 of 12 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	G	3	NGC	O10-C10-C11-O11
5	I	2	NGC	O10-C10-C11-O11
5	K	2	NGC	O10-C10-C11-O11
6	L	1	NAG	C4-C5-C6-O6
6	L	1	NAG	O5-C5-C6-O6



There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	I	2	NGC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	NAG	E	401	1	14,14,15	0.45	0	17,19,21	1.49	2 (11%)
7	NAG	A	401	1	14,14,15	0.39	0	17,19,21	1.12	1 (5%)
7	NAG	C	401	1	14,14,15	0.41	0	17,19,21	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	E	401	1	-	2/6/23/26	0/1/1/1
7	NAG	A	401	1	-	2/6/23/26	0/1/1/1
7	NAG	C	401	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	E	401	NAG	C1-O5-C5	4.16	117.83	112.19
7	A	401	NAG	C1-O5-C5	3.95	117.55	112.19
7	E	401	NAG	O5-C5-C6	3.01	111.93	107.20

There are no chirality outliers.



5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	E	401	NAG	O5-C5-C6-O6
7	E	401	NAG	C4-C5-C6-O6
7	A	401	NAG	O5-C5-C6-O6
7	A	401	NAG	C4-C5-C6-O6
7	C	401	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/334 (94%)	0.03	5 (1%) 72 74	20, 32, 56, 104	0
1	C	315/334 (94%)	-0.02	3 (0%) 82 84	21, 34, 63, 117	0
1	E	315/334 (94%)	2.24	123 (39%) 0 0	37, 74, 192, 260	0
2	B	170/186 (91%)	0.68	14 (8%) 11 12	23, 54, 80, 91	0
2	D	170/186 (91%)	1.24	40 (23%) 0 0	20, 66, 107, 119	0
2	F	170/186 (91%)	7.38	144 (84%) 0 0	55, 197, 250, 304	0
All	All	1455/1560 (93%)	1.57	329 (22%) 0 0	20, 51, 207, 304	0

The worst 5 of 329 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	152	ILE	28.2
2	F	118	LEU	28.0
2	F	29	ALA	25.4
2	F	119	TYR	22.0
2	F	45	ILE	21.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GAL	I	1	12/12	0.70	0.23	56,72,77,79	0
5	GAL	K	1	12/12	0.73	0.25	109,112,118,118	0
4	BMA	H	3	11/12	0.75	0.26	78,83,88,88	0
6	NAG	L	2	14/15	0.83	0.20	90,93,94,94	0
6	NAG	J	2	14/15	0.84	0.16	60,62,69,69	0
5	NGC	K	2	21/22	0.84	0.29	87,99,103,105	0
4	NAG	H	2	14/15	0.88	0.13	54,57,62,71	0
6	NAG	L	1	14/15	0.89	0.11	68,71,77,82	0
4	NAG	H	1	14/15	0.93	0.11	36,40,47,49	0
5	NGC	I	2	21/22	0.94	0.11	39,41,45,48	0
6	NAG	J	1	14/15	0.95	0.08	40,45,51,52	0
3	NGC	G	3	21/22	0.97	0.08	24,28,31,35	0
3	NAG	G	1	15/15	0.97	0.08	28,34,38,38	0
3	GAL	G	2	11/12	0.97	0.09	24,25,26,27	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain I:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain K:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain J:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	NAG	E	401	14/15	0.68	0.42	88,93,108,110	0
7	NAG	A	401	14/15	0.82	0.19	61,64,80,80	0
7	NAG	C	401	14/15	0.83	0.24	74,78,83,87	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.