

Full wwPDB X-ray Structure Validation Report (i)

Jul 20, 2022 – 02:24 pm BST

PDB ID : 8AB3

Title : Crystal Structure of the Lactate Dehydrogenase of Cyanobacterium Aponinum

in complex with oxamate, NADH and FBP.

Authors: Robin, A.Y.; Girard, E.; Madern, D.

Deposited on : 2022-07-04

Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.29

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0267$

CCP4 : 7.1.010 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.29

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	3797 (2.64-2.60)
Clashscore	141614	4168 (2.64-2.60)
Ramachandran outliers	138981	4093 (2.64-2.60)
Sidechain outliers	138945	4093 (2.64-2.60)
RSRZ outliers	127900	3731 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	337	91%	6% •
1	В	337	86%	11% •••
1	С	337	91%	7% •
1	D	337	92%	5% •



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10654 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called L-lactate dehydrogenase.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	Λ	330	Total	С	N	О	S	0	0	0
1	A	990	2498	1589	410	486	13	U	0	
1	В	328	Total	С	N	О	S	0	1	0
1	Б	320	2459	1567	401	478	13	0		
1	C	330	Total	С	N	О	S	0	1	0
1		330	2491	1587	411	480	13	U	1	
1	D	290	Total	С	N	О	S	0	1	0
		328	2475	1578	406	478	13	U	1	

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	334	HIS	-	expression tag	UNP K9Z684
A	335	HIS	-	expression tag	UNP K9Z684
A	336	HIS	-	expression tag	UNP K9Z684
A	337	HIS	-	expression tag	UNP K9Z684
A	338	HIS	-	expression tag	UNP K9Z684
A	339	HIS	-	expression tag	UNP K9Z684
В	334	HIS	-	expression tag	UNP K9Z684
В	335	HIS	-	expression tag	UNP K9Z684
В	336	HIS	-	expression tag	UNP K9Z684
В	337	HIS	-	expression tag	UNP K9Z684
В	338	HIS	-	expression tag	UNP K9Z684
В	339	HIS	_	expression tag	UNP K9Z684
С	334	HIS	-	expression tag	UNP K9Z684
С	335	HIS	_	expression tag	UNP K9Z684
С	336	HIS	-	expression tag	UNP K9Z684
С	337	HIS	-	expression tag	UNP K9Z684
С	338	HIS	-	expression tag	UNP K9Z684
С	339	HIS	-	expression tag	UNP K9Z684
D	334	HIS	-	expression tag	UNP K9Z684
D	335	HIS	-	expression tag	UNP K9Z684
D	336	HIS	-	expression tag	UNP K9Z684



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	337	HIS	-	expression tag	UNP K9Z684
D	338	HIS	-	expression tag	UNP K9Z684
D	339	HIS	-	expression tag	UNP K9Z684


• Molecule 2 is OXAMIC ACID (three-letter code: OXM) (formula: $C_2H_3NO_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C N O 6 2 1 3	0	0
2	В	1	Total C N O 6 2 1 3	0	0
2	С	1	Total C N O 6 2 1 3	0	0
2	D	1	Total C N O 6 2 1 3	0	0


• Molecule 3 is 1,6-di-O-phosphono-beta-D-fructofuranose (three-letter code: FBP) (formula: $C_6H_{14}O_{12}P_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	А	1	1 Total C O F		Р	0	0		
		1	20	6	12	2	O		
9	C	1	Total	\mathbf{C}	Ο	Р	0	0	
3		1	20	6	12	2	U		

• Molecule 4 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula: $C_{21}H_{29}N_7O_{14}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total 44	C 21	N 7	O 14	P 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
4	D	1	Total	С	N	О	Р	0	0	
4	В	1	44	21	7	14	2	U	0	
1	C	1	Total	С	N	О	Р	0	0	
4		1	44	21	7	14	2	U		
1	D	1	Total	С	N	О	Р	0	0	
4	4 D	1	44	21	7	14	2	U		

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	128	Total O 128 128	0	0
5	В	108	Total O 108 108	0	0
5	С	147	Total O 147 147	0	0
5	D	108	Total O 108 108	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: L-lactate dehydrogenase





4 Data and refinement statistics (i)

Property	Value	Sour				
Space group	P 1 21 1	Depos				
Cell constants	70.37Å 111.50Å 101.79Å	Depos				
a, b, c, α , β , γ	90.00° 99.78° 90.00°					
Resolution (Å)	100.31 - 2.62	Depos				
Resolution (A)	100.31 - 2.62	ED				
% Data completeness	99.3 (100.31-2.62)	Depos				
(in resolution range)	99.3 (100.31-2.62)	ED				
R_{merge}	0.19	Depos				
R_{sym}	(Not available)	Depos				
$< I/\sigma(I) > 1$	2.24 (at 2.62Å)	Xtria				
Refinement program	BUSTER 2.10.4 (8-JUN-2022), BUSTER 2.10.4 (8-JUN-2022)	Depos				
D D	0.186 , 0.232	Depos				
R, R_{free}	0.178 , 0.223	DC				
R_{free} test set	2317 reflections (4.98%)	wwPDI				
Wilson B-factor (Å ²)	35.5	Xtria				
Anisotropy	0.265	Xtria				
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	(Not available), (Not available)	ED				
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.31$	Xtria				
Estimated twinning fraction	No twinning to report.	Xtria				
F_o, F_c correlation	0.94	ED				
Total number of atoms	10654	wwPDI				
Average B, all atoms (Å ²)	39.0	wwPDI				

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.23% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAI, OXM, FBP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	$\mid RMSZ \mid \# Z > 5$		RMSZ	# Z > 5	
1	A	0.35	0/2535	0.54	0/3441	
1	В	0.37	0/2496	0.56	0/3395	
1	С	0.35	0/2528	0.53	0/3433	
1	D	0.34	0/2512	0.52	0/3412	
All	All	0.35	0/10071	0.54	0/13681	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2498	0	2539	11	0
1	В	2459	0	2471	18	0
1	С	2491	0	2536	15	0
1	D	2475	0	2517	11	0
2	A	6	0	2	0	0
2	В	6	0	2	0	0
2	С	6	0	2	0	0
2	D	6	0	2	0	0
3	A	20	0	10	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	С	20	0	10	0	0
4	A	44	0	27	0	0
4	В	44	0	27	1	0
4	С	44	0	27	0	0
4	D	44	0	27	1	0
5	A	128	0	0	1	0
5	В	108	0	0	0	0
5	С	147	0	0	2	0
5	D	108	0	0	0	0
All	All	10654	0	10199	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:297:SER:HB3	1:B:302:ILE:HD13	1.79	0.63
1:B:168:ASP:OD2	1:B:195:HIS:ND1	2.33	0.61
1:A:28:ILE:HD11	1:A:84:ILE:HD11	1.86	0.57
1:C:270:GLU:HB3	1:D:182:ILE:HA	1.87	0.56
1:B:33:VAL:HG21	4:B:402:NAI:C6N	2.37	0.55
1:B:302:ILE:HG22	1:B:303:LYS:HG3	1.90	0.53
1:C:28:ILE:HD11	1:C:84:ILE:HD11	1.89	0.53
1:B:218:TRP:O	1:B:221:LEU:HB2	2.11	0.51
1:A:270:GLU:HB3	1:B:182:ILE:HA	1.93	0.51
1:D:138:VAL:O	4:D:402:NAI:H2N	2.11	0.50
1:D:299:ARG:HB2	1:D:302:ILE:CD1	2.42	0.49
1:D:299:ARG:HB2	1:D:302:ILE:HD11	1.95	0.49
1:A:23:ARG:NH2	1:C:23:ARG:HG3	2.28	0.49
1:A:100:VAL:HG23	1:A:111:LEU:HD22	1.95	0.48
1:B:244:ILE:HG12	1:B:249:TYR:HA	1.94	0.47
1:C:100:VAL:HG23	1:C:111:LEU:HD22	1.95	0.47
1:B:222:PRO:HD2	1:B:225:GLU:HB2	1.97	0.47
1:D:328:PHE:O	1:D:331:ILE:HG13	2.15	0.47
1:C:121:LYS:O	5:C:501:HOH:O	2.21	0.46
1:B:30:LEU:HD21	1:B:52:LEU:HD22	1.98	0.46
1:C:140:ASN:HA	1:C:142:VAL:N	2.31	0.46
1:D:102:GLN:HB3	1:D:111:LEU:HD11	1.96	0.46
1:C:30:LEU:HD21	1:C:52:LEU:HD22	1.98	0.46
1:D:194:GLU:HG3	1:D:324:LEU:HD21	1.97	0.45



Continued from previous page...

A + 1		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:D:294:ARG:HG2	1:D:304:THR:HG23	1.99	0.45
1:B:89:ARG:HG2	1:B:129:TYR:HB3	1.99	0.45
1:B:109:LEU:HA	1:B:141:PRO:HG3	1.97	0.45
1:C:187:VAL:HA	1:C:208:VAL:HG12	1.99	0.45
1:A:178:LYS:HE2	5:A:545:HOH:O	2.16	0.45
1:A:294:ARG:HG2	1:A:304:THR:HG23	1.99	0.44
1:C:182:ILE:HA	1:D:270:GLU:HB3	1.97	0.44
1:B:140:ASN:HA	1:B:142:VAL:N	2.32	0.44
1:A:187:VAL:HA	1:A:208:VAL:HG12	2.00	0.44
1:C:294:ARG:HG2	1:C:304:THR:HG23	1.99	0.43
1:A:16:ASN:HA	1:A:17:PRO:HD3	1.91	0.43
1:B:314:GLU:O	1:B:318:GLN:HG2	2.18	0.43
1:C:114:ARG:HD3	5:C:571:HOH:O	2.19	0.43
1:D:187:VAL:HA	1:D:208:VAL:HG12	2.00	0.43
1:C:30:LEU:CD1	1:C:52:LEU:HB3	2.49	0.42
1:B:161:ILE:HD11	1:B:300:GLY:HA2	2.02	0.42
1:B:29:GLY:O	1:B:97:THR:OG1	2.37	0.42
1:A:20:LEU:HD23	1:C:298:GLU:HB2	2.01	0.41
1:A:30:LEU:HD13	1:A:52:LEU:HB3	2.02	0.41
1:C:30:LEU:HD13	1:C:52:LEU:HB3	2.01	0.41
1:B:52:LEU:HD11	1:B:79:ILE:HD11	2.03	0.41
1:B:59:LYS:O	1:B:63:GLU:HG2	2.19	0.41
1:B:270:GLU:HG3	1:B:297:SER:HB2	2.03	0.41
1:D:23:ARG:HD3	1:D:266:LEU:HD22	2.02	0.41
1:A:24:LYS:HG3	1:A:49:GLU:HG3	2.02	0.40
1:C:16:ASN:HA	1:C:17:PRO:HD3	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	328/337~(97%)	317 (97%)	11 (3%)	0	100 100
1	В	327/337~(97%)	315 (96%)	12 (4%)	0	100 100
1	С	329/337~(98%)	319 (97%)	10 (3%)	0	100 100
1	D	327/337~(97%)	316 (97%)	11 (3%)	0	100 100
All	All	$1311/1348 \; (97\%)$	1267 (97%)	44 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	274/288~(95%)	269 (98%)	5 (2%)	59 79		
1	В	265/288~(92%)	255 (96%)	10 (4%)	33 57		
1	\mathbf{C}	$272/288 \ (94\%)$	268 (98%)	4 (2%)	65 82		
1	D	$270/288 \ (94\%)$	269 (100%)	1 (0%)	91 97		
All	All	$1081/1152\ (94\%)$	1061 (98%)	20 (2%)	57 78		

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	LEU
1	A	86	ASN
1	A	142	VAL
1	A	223	GLU
1	A	286	GLU
1	В	20	LEU
1	В	30	LEU
1	В	49	GLU
1	В	113	GLU
1	В	125	ASP
1	В	178	LYS
1	В	208	VAL
1	В	216	ASN



Continued from previous page...

Mol	Chain	Res	Type
1	В	221	LEU
1	В	292	ILE
1	С	30	LEU
1	С	250	THR
1	С	286	GLU
1	С	302	ILE
1	D	250	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	102	GLN
1	A	318	GLN
1	С	102	GLN
1	С	318	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FBP	С	403	-	18,20,20	0.71	0	23,32,32	0.69	0
4	NAI	В	402	-	42,48,48	0.48	0	47,73,73	0.67	1 (2%)
4	NAI	D	402	-	42,48,48	0.50	0	47,73,73	0.73	1 (2%)
2	OXM	С	401	-	5,5,5	0.26	0	4,6,6	0.54	0
2	OXM	В	401	-	5,5,5	2.04	2 (40%)	4,6,6	1.88	2 (50%)
4	NAI	A	403	-	42,48,48	0.52	0	47,73,73	0.68	1 (2%)
2	OXM	A	401	-	5,5,5	1.99	2 (40%)	4,6,6	2.20	2 (50%)
2	OXM	D	401	-	5,5,5	0.21	0	4,6,6	1.13	0
4	NAI	С	402	-	42,48,48	0.50	1 (2%)	47,73,73	0.76	1 (2%)
3	FBP	A	402	-	18,20,20	0.62	0	23,32,32	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FBP	С	403	-	-	1/13/32/32	0/1/1/1
4	NAI	В	402	-	-	11/25/72/72	0/5/5/5
4	NAI	D	402	-	-	17/25/72/72	0/5/5/5
2	OXM	С	401	-	-	0/3/4/4	-
2	OXM	В	401	_	ı	0/3/4/4	-
4	NAI	A	403	_	-	11/25/72/72	0/5/5/5
2	OXM	A	401	-	-	0/3/4/4	-
2	OXM	D	401	-	-	0/3/4/4	-
4	NAI	С	402	_	-	5/25/72/72	0/5/5/5
3	FBP	A	402	_	-	3/13/32/32	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	A	401	OXM	O2-C2	3.57	1.32	1.22
2	В	401	OXM	O2-C2	3.49	1.31	1.22
2	В	401	OXM	O3-C2	-2.88	1.22	1.30
2	A	401	OXM	O3-C2	-2.58	1.23	1.30
4	С	402	NAI	C8A-N7A	-2.04	1.31	1.34

All (8) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	401	OXM	O3-C2-C1	3.33	121.55	113.84
2	В	401	OXM	O3-C2-C1	2.94	120.64	113.84
2	A	401	OXM	O2-C2-C1	-2.83	115.76	122.06
4	D	402	NAI	C5A-C6A-N6A	2.44	124.06	120.35
4	В	402	NAI	C5A-C6A-N6A	2.25	123.78	120.35
2	В	401	OXM	O2-C2-C1	-2.24	117.09	122.06
4	С	402	NAI	C5A-C6A-N6A	2.21	123.71	120.35
4	A	403	NAI	C5A-C6A-N6A	2.17	123.66	120.35

There are no chirality outliers.

All (48) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	FBP	C1-O1-P1-O1P
4	A	403	NAI	C5B-O5B-PA-O1A
4	В	402	NAI	C5B-O5B-PA-O1A
4	В	402	NAI	C2N-C3N-C7N-O7N
4	D	402	NAI	C5B-O5B-PA-O1A
4	D	402	NAI	PN-O3-PA-O5B
4	D	402	NAI	C5D-O5D-PN-O2N
4	В	402	NAI	O4B-C4B-C5B-O5B
4	D	402	NAI	O4D-C4D-C5D-O5D
4	D	402	NAI	C3D-C4D-C5D-O5D
4	D	402	NAI	O4B-C4B-C5B-O5B
4	D	402	NAI	C2D-C1D-N1N-C2N
4	A	403	NAI	PN-O3-PA-O5B
4	В	402	NAI	PN-O3-PA-O5B
4	A	403	NAI	O4B-C4B-C5B-O5B
3	A	402	FBP	C1-O1-P1-O2P
4	A	403	NAI	C5B-O5B-PA-O3
4	В	402	NAI	C5B-O5B-PA-O3
4	D	402	NAI	C5B-O5B-PA-O3
4	A	403	NAI	PA-O3-PN-O2N
4	D	402	NAI	PA-O3-PN-O2N
4	D	402	NAI	O4D-C1D-N1N-C2N
4	A	403	NAI	C5B-O5B-PA-O2A
4	В	402	NAI	C5B-O5B-PA-O2A
4	В	402	NAI	C2N-C3N-C7N-N7N
4	D	402	NAI	C5B-O5B-PA-O2A
4	D	402	NAI	C5D-O5D-PN-O1N
4	D	402	NAI	C2D-C1D-N1N-C6N
4	A	403	NAI	O4D-C1D-N1N-C2N
4	D	402	NAI	O4D-C1D-N1N-C6N



Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	С	402	NAI	O4D-C1D-N1N-C2N
4	A	403	NAI	C2D-C1D-N1N-C2N
4	С	402	NAI	C2D-C1D-N1N-C2N
4	В	402	NAI	C3B-C4B-C5B-O5B
4	В	402	NAI	O4D-C1D-N1N-C2N
4	С	402	NAI	O4B-C4B-C5B-O5B
3	A	402	FBP	O1-C1-C2-C3
3	С	403	FBP	O1-C1-C2-C3
4	D	402	NAI	C3B-C4B-C5B-O5B
4	С	402	NAI	O4D-C1D-N1N-C6N
4	A	403	NAI	O4D-C1D-N1N-C6N
4	D	402	NAI	C5D-O5D-PN-O3
4	С	402	NAI	C2D-C1D-N1N-C6N
4	A	403	NAI	PA-O3-PN-O1N
4	В	402	NAI	PA-O3-PN-O1N
4	D	402	NAI	PA-O3-PN-O1N
4	A	403	NAI	C2D-C1D-N1N-C6N
4	В	402	NAI	C2D-C1D-N1N-C2N


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	402	NAI	1	0
4	D	402	NAI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(A^2)$	Q<0.9
1	A	330/337 (97%)	-0.35	0 100 100	22, 36, 59, 74	0
1	В	328/337 (97%)	-0.26	6 (1%) 68 64	24, 40, 66, 79	0
1	С	330/337 (97%)	-0.32	2 (0%) 89 88	23, 34, 58, 70	0
1	D	328/337 (97%)	-0.34	0 100 100	23, 38, 60, 72	0
All	All	1316/1348 (97%)	-0.32	8 (0%) 89 88	22, 37, 62, 79	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	107	SER	4.1
1	В	110	SER	2.8
1	В	105	GLY	2.8
1	В	4	PHE	2.5
1	С	3	MET	2.5
1	С	17	PRO	2.3
1	В	241	TYR	2.3
1	В	109	LEU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	OXM	D	401	6/6	0.90	0.14	44,45,45,46	0
2	OXM	С	401	6/6	0.95	0.14	41,42,42,43	0
2	OXM	В	401	6/6	0.96	0.18	44,44,45,45	0
3	FBP	С	403	20/20	0.96	0.18	37,38,39,40	0
4	NAI	В	402	44/44	0.96	0.16	49,50,53,53	0
4	NAI	D	402	44/44	0.96	0.14	38,43,48,48	0
4	NAI	A	403	44/44	0.97	0.13	28,30,32,33	0
3	FBP	A	402	20/20	0.97	0.15	39,41,42,42	0
2	OXM	A	401	6/6	0.97	0.15	40,40,41,41	0
4	NAI	С	402	44/44	0.98	0.14	27,28,29,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around OXM D 401: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)








Electron density around OXM B 401: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAI B 402:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAI D 402:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAI A 403:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around FBP A 402:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















6.5 Other polymers (i)

There are no such residues in this entry.

