



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 11, 2024 – 02:06 PM EDT

PDB ID : 8S9L
Title : Structure of monomeric FAM111A SPD V347D Mutant
Authors : Palani, S.; Alvey, J.A.; Cong, A.T.Q.; Schellenberg, M.J.; Machida, Y.
Deposited on : 2023-03-29
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	
1	B	270	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3894 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine protease FAM111A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	241	1935	1251	320	355	9	0	1	0
1	B	233	1864	1205	309	341	9	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	342	GLY	-	expression tag	UNP Q96PZ2
A	343	HIS	-	expression tag	UNP Q96PZ2
A	344	MET	-	expression tag	UNP Q96PZ2
A	347	ASP	VAL	engineered mutation	UNP Q96PZ2
B	342	GLY	-	expression tag	UNP Q96PZ2
B	343	HIS	-	expression tag	UNP Q96PZ2
B	344	MET	-	expression tag	UNP Q96PZ2
B	347	ASP	VAL	engineered mutation	UNP Q96PZ2

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	O S	0	0
			5	4 1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	52	Total	O	0	0
			52	52		
3	B	38	Total	O	0	0
			38	38		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine protease FAM111A



- Molecule 1: Serine protease FAM111A



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.46Å 70.03Å 128.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.96 – 1.85 48.96 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.96-1.85) 99.8 (48.96-1.85)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.86Å)	Xtrriage
Refinement program	PHENIX (1.17.1_3660: ???)	Depositor
R, R_{free}	0.218 , 0.252 0.218 , 0.252	Depositor DCC
R_{free} test set	2003 reflections (3.77%)	wwPDB-VP
Wilson B-factor (Å ²)	39.3	Xtrriage
Anisotropy	0.472	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 59.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.034 for k,h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3894	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/1987	0.68	0/2688
1	B	0.59	0/1909	0.66	0/2582
All	All	0.59	0/3896	0.67	0/5270

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1935	0	1900	21	0
1	B	1864	0	1832	17	0
2	B	5	0	0	0	0
3	A	52	0	0	2	0
3	B	38	0	0	0	0
All	All	3894	0	3732	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:481:GLN:NE2	3:A:703:HOH:O	2.32	0.63
1:B:458:ASN:HD21	1:B:552:LEU:H	1.45	0.62
1:A:343:HIS:CE1	1:A:345:LYS:HB3	2.39	0.58
1:A:490:GLN:HG2	1:A:493:ARG:CZ	2.34	0.57
1:B:524:VAL:HG12	1:B:525:HIS:ND1	2.20	0.57
1:B:498:GLN:O	1:B:502:GLN:HG2	2.06	0.55
1:B:496:LYS:O	1:B:500:ARG:HG2	2.08	0.54
1:A:396:GLU:HB2	1:A:399:LYS:HE2	1.91	0.53
1:B:423:ASN:ND2	1:B:450:GLN:HE22	2.05	0.53
1:B:423:ASN:HD22	1:B:450:GLN:HE22	1.57	0.53
1:B:496:LYS:HD2	1:B:500:ARG:CZ	2.40	0.52
1:A:526:ASN:OD1	1:A:528:ASP:HB2	2.11	0.51
1:A:471:ILE:HG12	1:A:545:VAL:HG22	1.94	0.50
1:B:518:ARG:HE	1:B:521:GLN:HB3	1.77	0.49
1:A:411:THR:HG22	1:A:413:GLY:H	1.78	0.47
1:B:590:LYS:HE2	1:B:594:GLU:OE2	2.15	0.46
1:A:496:LYS:O	1:A:500:ARG:HG3	2.15	0.46
1:B:496:LYS:HD2	1:B:500:ARG:NH2	2.30	0.45
1:B:498:GLN:HB3	1:B:517:GLN:OE1	2.16	0.45
1:B:384:ARG:HG2	1:B:436:GLU:HA	1.98	0.45
1:A:590:LYS:O	1:A:594:GLU:HG3	2.17	0.45
1:A:493:ARG:NH2	1:A:529:VAL:O	2.49	0.44
1:A:428:GLU:CD	1:A:446:LYS:HG2	2.39	0.43
1:B:479:LYS:CG	1:B:480:LYS:H	2.31	0.43
1:A:461:THR:CG2	1:A:553:VAL:HG12	2.48	0.43
1:A:409:ARG:HG3	3:A:750:HOH:O	2.20	0.42
1:A:472:ILE:O	1:A:544:PRO:HD2	2.20	0.42
1:A:343:HIS:HB3	1:A:346:VAL:HG12	2.00	0.42
1:A:401:ALA:HB1	1:A:429:PRO:C	2.40	0.42
1:A:531:THR:HG22	1:A:574:PHE:HB3	2.02	0.41
1:A:461:THR:HG21	1:A:553:VAL:CG1	2.51	0.41
1:B:363:ASP:HA	1:B:367:THR:HG22	2.02	0.41
1:A:435:ASN:HB3	1:A:440:TYR:CE2	2.56	0.41
1:B:518:ARG:HH21	1:B:521:GLN:HB3	1.85	0.41
1:B:461:THR:HG21	1:B:551:SER:HB2	2.02	0.41
1:A:401:ALA:HB1	1:A:429:PRO:O	2.21	0.41
1:A:396:GLU:OE1	1:A:399:LYS:NZ	2.38	0.40
1:B:586:LYS:HG2	1:B:593:TYR:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/270 (87%)	225 (95%)	11 (5%)	0	100	100
1	B	225/270 (83%)	217 (96%)	8 (4%)	0	100	100
All	All	461/540 (85%)	442 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/238 (89%)	210 (100%)	1 (0%)	88	86
1	B	204/238 (86%)	200 (98%)	4 (2%)	55	40
All	All	415/476 (87%)	410 (99%)	5 (1%)	71	62

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	345	LYS
1	B	423	ASN
1	B	481	GLN
1	B	500	ARG
1	B	518	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	343	HIS
1	A	599	ASN
1	B	423	ASN
1	B	458	ASN
1	B	474	HIS
1	B	490	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	701	-	4,4,4	0.14	0	6,6,6	0.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/270 (89%)	0.94	33 (13%) 3 3	32, 62, 105, 132	0
1	B	233/270 (86%)	0.90	32 (13%) 3 3	37, 65, 107, 133	0
All	All	474/540 (87%)	0.92	65 (13%) 3 3	32, 63, 106, 133	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	516	THR	5.7
1	B	344	MET	4.5
1	B	564	TYR	4.2
1	B	394	GLY	4.2
1	B	598	VAL	4.0
1	A	343	HIS	4.0
1	A	344	MET	4.0
1	A	396	GLU	4.0
1	B	479	LYS	4.0
1	B	521	GLN	3.9
1	B	413	GLY	3.8
1	A	394	GLY	3.8
1	B	399	LYS	3.8
1	A	430	TRP	3.7
1	B	396	GLU	3.7
1	A	393	ASP	3.6
1	A	516	THR	3.6
1	A	399	LYS	3.5
1	B	393	ASP	3.4
1	A	592	TRP	3.3
1	B	518	ARG	3.3
1	A	524	VAL	3.3
1	B	502	GLN	3.3
1	A	521	GLN	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	497	CYS	3.2
1	A	520	PHE	3.1
1	A	397	PRO	3.1
1	A	518	ARG	3.1
1	A	507	GLU	3.0
1	B	600	GLN	3.0
1	B	499	GLU	3.0
1	A	596	VAL	3.0
1	A	395	ILE	2.9
1	A	391	VAL	2.9
1	A	495	LYS	2.9
1	B	501	VAL	2.8
1	B	495	LYS	2.7
1	B	520	PHE	2.7
1	B	397	PRO	2.6
1	B	474	HIS	2.6
1	A	506	ALA	2.6
1	A	564	TYR	2.5
1	A	398	SER	2.5
1	A	504	LYS	2.5
1	A	422	THR	2.5
1	B	463	VAL	2.5
1	A	599	ASN	2.3
1	B	517	GLN	2.3
1	B	595	GLU	2.3
1	B	537	PHE	2.2
1	A	429	PRO	2.2
1	A	560	PHE	2.2
1	B	432	GLU	2.2
1	A	464	PRO	2.2
1	A	423	ASN	2.1
1	B	494	ALA	2.1
1	B	395	ILE	2.1
1	B	352	ARG	2.1
1	B	599	ASN	2.1
1	A	588	ARG	2.1
1	A	517	GLN	2.1
1	A	392	GLY	2.1
1	B	593	TYR	2.0
1	A	553	VAL	2.0
1	B	433	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	701	5/5	0.86	0.15	132,134,135,135	0

6.5 Other polymers [i](#)

There are no such residues in this entry.